arXiv:1203.3187v2 [math.CO] 12 Sep 2012

MULTIPLY-REFINED ENUMERATION OF ALTERNATING SIGN MATRICES

ROGER E. BEHREND

ABSTRACT. Four natural boundary statistics and two natural bulk statistics are considered for alternating sign matrices (ASMs). Specifically, these statistics are the positions of the 1's in the first and last row and column of an ASM, and the numbers of generalized inversions and -1's in an ASM. Previously-known results for the exact enumeration of ASMs with prescribed values of some of these statistics are reviewed in detail. A quadratic relation which fully determines the generating function associated with all six statistics is then obtained. The derivation of the relation involves combining the Desnanot–Jacobi determinant identity with the Izergin–Korepin formula for the partition function of the six-vertex model with domain-wall boundary conditions.

CONTENTS

1. Introduction 2
2. Definitions and Basic Properties 4
2.1. Statistics 4
2.2. Generating functions 7
3. Previously-known and Related Results 9
3.1. Bulk parameter $y = 0$ 9
3.2. Bulk parameters satisfying $y = x + 1$ 10
3.3. Bulk parameters $x = y = 1$ 13
3.4. Arbitrary bulk parameters x and y 19
3.5. ASM enumeration involving statistics associated with several rows or several columns 21
3.6. ASMs with several rows or columns closest to two opposite boundaries prescribed 24
3.7. The cases $Z_n(1, 3)$ and $Z_n(1, 3; z)$ 25
3.8. The case $Z_n(1, 1; -1)$ 25
3.9. ASMs with a fixed number of generalized inversions 25
3.10. ASMs with a fixed number of -1's 25
3.11. Objects in simple bijection with ASMs 26
3.12. Descending plane partitions 26
3.13. Totally symmetric self-complementary plane partitions 27
3.14. Loop models 28
3.15. ASMs invariant under symmetry operations 28
4. New Results 29
4.1. Main result 29

Key words and phrases. Alternating sign matrices, six-vertex model with domain-wall boundary conditions, Desnanot–Jacobi identity.

4.2. Consequences for arbitrary bulk parameters x and y 30
4.3. Consequences for bulk parameters $x = y = 1$ 34
5. Proofs 35
5.1. A generalized ASM generating function 35
5.2. The bijection between ASMs and configurations of the six-vertex model with DWBC 37
5.3. The partition function of the six-vertex model with DWBC 40
5.4. The Izergin–Korepin determinant formula 41
5.5. The Desnanot–Jacobi identity 42
5.6. Proof of Theorem 1 43
5.7. Alternative statement of some results of Section 4.1 44
5.8. Derivations of (19) and (20) 47
5.9. Derivation of (71) 48
5.10. Derivations of (48) and (73) 49
References 50

1. INTRODUCTION

A major focus of attention throughout the history of alternating sign matrices (ASMs) has simply been the derivation of results related to their exact enumeration. Such results typically state that the number of ASMs which satisfy specific conditions, such as having prescribed values of certain statistics or being invariant under certain symmetry operations, is given by an explicit formula or generating function, is equal to the number of combinatorial objects of some other variety satisfying specific conditions, or is equal to a number which arises from a particular physical model.

A few examples of such results, with references to conjectures and initial proofs, are as follows: a formula for the total number of ASMs of arbitrary fixed size but with no further conditions applied (Mills, Robbins and Rumsey [102, 103], Zeilberger [145], and Kuperberg [92]); a formula for the number of ASMs with a prescribed boundary row or column (Mills, Robbins and Rumsey [102, 103], and Zeilberger [146]); formulae for numbers of ASMs invariant under certain natural symmetry operations (Robbins [123, 124], Kuperberg [93], Okada [111], and Razumov and Stroganov [119, 120]); equalities between numbers of certain ASMs and numbers of certain totally symmetric self-complementary plane partitions (Mills, Robbins and Rumsey [104], and Fonseca and Zinn-Justin [75]); equalities between numbers of certain ASMs and numbers of certain descending plane partitions (Mills, Robbins and Rumsey [103], and Behrend, Di Francesco and Zinn-Justin [17]); equalities between numbers of certain ASMs and numbers associated with a certain case of the $O(1)$ loop model (Razumov and Stroganov [116], and Cantini and Sportiello [31]). For reviews of some of these results, see, for example, Bressoud [22, 24], Bressoud and Propp [25], Hone [78], Zeilberger [148], or Zinn-Justin [149].

The main result in this paper is of the type which provides an expression for the generating function associated with numbers of ASMs of arbitrary fixed size with prescribed values of particular statistics. In any ASM, the first and last row and column each contain a single entry of 1, with all of their other entries being 0's, so that the positions of these 1's provide four natural statistics which describe the boundary configuration of an ASM. The generating function under primary consideration in this paper is associated with these four boundary statistics, together with two statistics which depend on the bulk structure of an ASM, namely the number of generalized inversions (which will be defined in (3)) and the number of -1 entries. Elementary properties of this generating function and these statistics will be outlined in Section 2.

All previously-known results for the exact enumeration of ASMs have involved fewer than six of these statistics. A review of such results will be given in Section 3. That section will aim to provide a comprehensive account of all previous appearances of these statistics in exact enumeration, together with an outline of some further related results which can be obtained straightforwardly from the previously-known results. Recently, a result involving all four of the boundary statistics, but neither of the bulk statistics, was obtained independently by Ayyer and Romik [12, Thm. 2]. That result will be reviewed in Section 3.3.

In ASM enumerations, it is common to refer to a certain order of refinement which is based on the number of boundary statistics involved. Hence, the primary generating functions used in this paper, and by Ayyer and Romik [12], can be described as quadruply-refined.

The main result of this paper, Theorem 1, will be stated in Section 4, and proved in Section 5. The approach used in the proof is essentially as follows. First, certain standard techniques, which have played a crucial role in proofs of many of the other known enumerative results for ASMs, are used to obtain a determinantal expression related to ASM generating functions. More specifically, a bijection between ASMs and configurations of the statistical mechanical six-vertex, or square ice, model with domain-wall boundary conditions (DWBC) is used to derive a relation between ASM generating functions and the partition function of the model, and the Izergin–Korepin formula is used to provide an explicit determinantal expression for this partition function. Next, the Desnanot–Jacobi determinant identity is applied to the matrix in the Izergin–Korepin formula, which leads to a quadratic relation involving the required quadruply-refined ASM generating function, and generating functions associated with no ASM boundary statistics or with two ASM boundary statistics, for all four pairs of adjacent boundaries. It will also be seen that this relation, together with simple initial conditions, fully determines the six-statistic quadruply-refined ASM generating function.

The key step in this derivation is the use of the Desnanot–Jacobi identity together with the Izergin–Korepin formula, and the reason that this leads to a relation of the previously-described form can be understood relatively easily. The Izergin–Korepin formula involves a certain prefactor, which depends on so-called spectral parameters $u_1, \ldots, u_n$ and $v_1, \ldots, v_n$, and the determinant of a certain $n \times n$ matrix, whose entry in row i and column j depends on u_i and v_j. Furthermore, in terms of generating functions for $n \times n$ ASMs, the parameters u_i and v_j are, in a certain sense, associated with row i and column j of the ASMs. For an $n \times n$ matrix M, the Desnanot–Jacobi identity (in the form used here) consists of a relation involving $\det M$ and five minors of M: the four connected $(n-1) \times (n-1)$ minors, corresponding to deletion of the first or last row and first or last column of M, and the central $(n-2)\times(n-2)$ minor, corresponding to deletion of the first and last row and column of M. Therefore, it seems that applying the Desnanot–Jacobi identity to the matrix in the Izergin–Korepin formula, and using an assignment of spectral parameters in which u_1, u_n, v_1 and v_n remain arbitrary, while $u_2,\ldots,u_{n-1}$ are equal and $v_2,\ldots,v_{n-1}$ are equal, should lead to an expression involving a quadruply-refined generating function for $n\times n$ ASMs, four doubly-refined generating functions for $(n-1)\times(n-1)$ ASMs, and an unrefined generating function for $(n-2)\times(n-2)$ ASMs. Indeed, this essentially does occur, with some further, subsidiary aspects of the resulting expression being related to the form of the prefactor in the Izergin–Korepin formula, and to the fact that each corner entry of an ASM is associated with two of the parameters u_1, u_n, v_1 or v_n.

A similar method, involving the combination of a different form of the Desnanot–Jacobi identity with the Izergin–Korepin formula, was used by Behrend, Di Francesco and Zinn-Justin [16], as part of the proof of a result concerning a doubly-refined enumeration of ASMs and descending plane partitions.

The Desnanot–Jacobi identity has, in fact, been associated with ASMs since these matrices first arose. In particular, the identity is used in Dodgson's condensation algorithm [60] for determinant evaluation, and it was through studies of this algorithm by Mills, Robbins and Rumsey that ASMs initially appeared. In particular, it was shown by Robbins and Rumsey [125, Eq. (27)] that if Dodgson's algorithm is applied to an $n\times n$ matrix M using a modified form of the Desnanot–Jacobi identity containing a parameter λ, then the resulting so-called λ-determinant of M can be expressed naturally as a sum over $n\times n$ ASMs. The latter expression involves the two ASM bulk statistics which are used in this paper, and, as will be shown in Section 3.2, the λ-determinant shares certain features with the quadruply-refined ASM generating function studied in this paper. For example, the modified Desnanot–Jacobi identity satisfied by the λ-determinant is analogous to the quadratic relation satisfied by the quadruply-refined ASM generating function. For further information regarding Dodgson condensation, λ-determinants, and related matters, see, for example, Abeles [1], Bressoud [22, Sec. 3.5], Bressoud and Propp [25], Propp [114], or Robbins and Rumsey [125].

Finally, it should be noted that the Desnanot–Jacobi identity, the λ-modified Desnanot–Jacobi identity, and further related equations studied by Robbins and Rumsey [125], and others, are closely connected with the Laurent phenomenon, cluster algebras, and associated areas. For reviews of such connections, see, for example, Hone [78, Sec. 5], or Propp [113, Sec. 10].

2. DEFINITIONS AND BASIC PROPERTIES

2.1. Statistics.

In this section, the standard definitions of ASMs and certain statistics for ASMs are given, and some simple properties of these matrices and statistics are identified.

An ASM, as first defined by Mills, Robbins and Rumsey [102, 103], is simply a square matrix in which each entry is 0, 1 or -1, and along each row and column the nonzero entries alternate in sign and have a sum of 1.

It follows that, for any ASM A,

$$\text{Each partial row sum } \textstyle\sum_{j'=1}^{j} A_{ij'} \text{ and each partial column sum } \sum_{i'=1}^{i} A_{i'j} \text{ is 0 or 1.} \qquad (1)$$

It can also be seen that any permutation matrix is an ASM, and that, in any ASM, the first and last row and column each contain a single 1, with all of their other entries being 0's.

For each positive integer n, denote the set of all $n \times n$ ASMs as $\mathrm{ASM}(n)$. For example, for $n = 1, 2, 3$, these sets are

$$\begin{aligned}
&\mathrm{ASM}(1) = \{(1)\},\\
&\mathrm{ASM}(2) = \left\{\begin{pmatrix}1&0\\0&1\end{pmatrix}, \begin{pmatrix}0&1\\1&0\end{pmatrix}\right\},\\
&\mathrm{ASM}(3) = \left\{\begin{pmatrix}1&0&0\\0&1&0\\0&0&1\end{pmatrix}, \begin{pmatrix}0&1&0\\1&0&0\\0&0&1\end{pmatrix}, \begin{pmatrix}1&0&0\\0&0&1\\0&1&0\end{pmatrix}, \begin{pmatrix}0&1&0\\0&0&1\\1&0&0\end{pmatrix}, \begin{pmatrix}0&0&1\\1&0&0\\0&1&0\end{pmatrix}, \begin{pmatrix}0&0&1\\0&1&0\\1&0&0\end{pmatrix}, \begin{pmatrix}0&1&0\\1&-1&1\\0&1&0\end{pmatrix}\right\}. \qquad (2)
\end{aligned}$$

For any $A \in \mathrm{ASM}(n)$, define statistics which depend on the bulk structure of A as

$$\nu(A) = \sum_{\substack{1 \le i < i' \le n\\ 1 \le j' \le j \le n}} A_{ij}\, A_{i'j'}, \qquad \mu(A) = \text{number of } -1\text{'s in } A, \qquad (3)$$

and define statistics which describe the configuration of A at its top, right, bottom and left boundaries as, respectively,

$$\begin{aligned}
\rho_{\mathrm{T}}(A) &= \text{number of 0's to the left of the 1 in the top row of } A,\\
\rho_{\mathrm{R}}(A) &= \text{number of 0's below the 1 in the right-most column of } A,\\
\rho_{\mathrm{B}}(A) &= \text{number of 0's to the right of the 1 in the bottom row of } A,\\
\rho_{\mathrm{L}}(A) &= \text{number of 0's above the 1 in the left-most column of } A. \qquad (4)
\end{aligned}$$

The statistics of (4) are given diagrammatically by

$$\begin{array}{ccc}
 & \leftarrow \rho_{\mathrm{T}}(A) \rightarrow \qquad\qquad & \\
\begin{array}{c}\uparrow\\ \rho_{\mathrm{L}}(A)\\ \downarrow\\ \\ \\ \\ \\ \\ \end{array} &
\begin{pmatrix}
0&0&0&0&1&0&0&0&0\\
0&&&&&&&&0\\
0&&&&&&&&0\\
1&&&&&&&&0\\
0&&&&A&&&&1\\
0&&&&&&&&0\\
0&&&&&&&&0\\
0&&&&&&&&0\\
0&0&1&0&0&0&0&0&0
\end{pmatrix} &
\begin{array}{c} \\ \\ \\ \\ \\ \uparrow\\ \rho_{\mathrm{R}}(A)\\ \downarrow\end{array} \\
 & \qquad\quad \longleftarrow \rho_{\mathrm{B}}(A) \longrightarrow &
\end{array} . \qquad (5)$$

The statistic $\nu(A)$ in (3) is a nonnegative integer for any $A \in \mathrm{ASM}(n)$, since it can be written as $\nu(A) = \sum_{i,j=1}^{n}(\sum_{i'=1}^{i-1} A_{i'j})(\sum_{j'=1}^{j} A_{ij'})$, and (1) implies that each factor in the summand is 0 or 1. This statistic can also be written as $\nu(A) = \sum_{1\le i\le i'\le n;\; 1\le j'<j\le n} A_{ij}\, A_{i'j'}$, where this can be obtained from the definition in (3) using the fact that each complete row and column sum of A is constant. As will be seen in Section 5.2, for the configuration of the six-vertex model with DWBC which corresponds to A, $\nu(A)$ is simply the number of vertex configurations of type (1), or equally the number of vertex configurations of type (2).

If A is a permutation matrix, then $\nu(A)$ is the number of inversions in the permutation π given by $\delta_{\pi_i,j} = A_{ij}$. Accordingly, for any ASM A, $\nu(A)$ is referred to as the number of generalized inversions in A. The statistic $\nu(A)$ was first defined and used by Robbins and Rumsey [125, Eq. (18)], who referred to it as the number of positive inversions in A [125, p. 182]. A closely-related statistic, $\sum_{1\le i<i'\le n;\; 1\le j'<j\le n} A_{ij}\, A_{i'j'} = \nu(A)+\mu(A)$, was previously defined and used by Mills, Robbins and Rumsey [103, p. 344], and provides an alternative generalized inversion number for each $A \in \mathrm{ASM}(n)$.

Certain generalizations of the statistics (3)–(4) will be defined in Section (3.5).

It can be checked easily that transposition or anticlockwise quarter-turn rotation of an ASM give another ASM, and that, for each $A \in \mathrm{ASM}(n)$, the statistics (3)–(4) behave under these operations according to

$$\begin{aligned}
&\nu(A) = \nu(A^T) = \tfrac{n(n-1)}{2} - \nu(A^Q) - \mu(A), && \mu(A) = \mu(A^T) = \mu(A^Q),\\
&\rho_{\mathrm{T}}(A) = \rho_{\mathrm{L}}(A^T) = n-1-\rho_{\mathrm{L}}(A^Q), && \rho_{\mathrm{R}}(A) = \rho_{\mathrm{B}}(A^T) = n-1-\rho_{\mathrm{T}}(A^Q),\\
&\rho_{\mathrm{B}}(A) = \rho_{\mathrm{R}}(A^T) = n-1-\rho_{\mathrm{R}}(A^Q), && \rho_{\mathrm{L}}(A) = \rho_{\mathrm{T}}(A^T) = n-1-\rho_{\mathrm{B}}(A^Q),
\end{aligned} \tag{6}$$

where Q denotes anticlockwise quarter-turn rotation, i.e., $A^Q_{ij} = A_{j,n+1-i}$. By combining transposition and anticlockwise quarter-turn rotation, the behavior of any of the eight operations of the dihedral group acting on ASMs can be obtained.

The properties of ASMs with a 1 in a corner can be described easily. For example, for the case of ASMs with a 1 in the top-left corner, the sets $\{A \in \mathrm{ASM}(n) \mid A_{11} = 1\} = \{A \in \mathrm{ASM}(n) \mid \rho_{\mathrm{T}}(A) = \rho_{\mathrm{L}}(A) = 0\}$ and $\mathrm{ASM}(n-1)$ are in bijection for any $n \ge 2$, where an ASM from the first set is mapped to the second set by simply deleting the first row and first column. Furthermore, if A from the first set is mapped to A' in the second set, then $\nu(A') = \nu(A)$, $\mu(A') = \mu(A)$, $\rho_{\mathrm{R}}(A') = \rho_{\mathrm{R}}(A)$ and $\rho_{\mathrm{B}}(A') = \rho_{\mathrm{B}}(A)$.

The properties of ASMs in which a 1 on a boundary is separated from a corner by a single zero, i.e., ASMs in which a boundary statistic of (4) is 1 or $n-2$, can also be described relatively easily. For example, for the case of ASMs A with $A_{21} = 1$, the sets $\{A \in \mathrm{ASM}(n) \mid A_{21} = A_{1,k+1} = 1\} = \{A \in \mathrm{ASM}(n) \mid \rho_{\mathrm{L}}(A) = 1,\ \rho_{\mathrm{T}}(A) = k\}$ and $\{A \in \mathrm{ASM}(n-1) \mid A_{11} = \ldots = A_{1,k-1} = 0\} = \{A \in \mathrm{ASM}(n-1) \mid \rho_{\mathrm{T}}(A) \ge k-1\}$ are in bijection for any $n \ge 2$ and $k = 1, \ldots, n-1$, where an ASM A from the first set is mapped to the second set by replacing $A_{2,k+1}$ by $A_{2,k+1}+1$, while leaving all other entries unchanged, and then deleting the first row and first column. Furthermore, if A from the first set is mapped to A' in the second set, then $\nu(A') = \nu(A) - A_{2,k+1} - 1$ and $\mu(A') = \mu(A) + A_{2,k+1}$.

2.2. **Generating functions.** In this section, various ASM generating functions are defined, and some simple relations which they satisfy are derived. Each of these generating functions will be labeled by a certain order of refinement, which corresponds to the number of boundary statistics of (4) (or simply the number of boundaries) with which it is associated. Each generating function is also associated with the two bulk statistics of (3), i.e., the numbers of generalized inversions and -1's in an ASM.

For each positive integer n, define a quadruply-refined ASM generating function, which involves all six statistics of (3)–(4), and associated indeterminates x, y, z_1, z_2, z_3 and z_4, as

$$Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{\rho_{\mathrm{R}}(A)}\,z_3^{\rho_{\mathrm{B}}(A)}\,z_4^{\rho_{\mathrm{L}}(A)}. \tag{7}$$

It follows that x and y can be regarded as bulk parameters or weights, and that z_1, z_2, z_3 and z_4 can be regarded as boundary parameters or weights.

Examples of the quadruply-refined ASM generating function, for $n=1,2,3$, are

$$\begin{aligned}
&Z_1^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)=1,\\
&Z_2^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)=1+x\,z_1\,z_2\,z_3\,z_4,\\
&Z_3^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)=1+x\,z_1\,z_4+x\,z_2\,z_3+x^2\,z_1\,z_2\,z_3^2\,z_4^2+x^2\,z_1^2\,z_2^2\,z_3\,z_4+\\
&\qquad\qquad x^3\,z_1^2\,z_2^2\,z_3^2\,z_4^2+x\,y\,z_1\,z_2\,z_3\,z_4,
\end{aligned} \tag{8}$$

where the terms are written in orders which correspond to those used in (2).

Now define triply-refined, adjacent-boundary doubly-refined, opposite-boundary doubly-refined, singly-refined and unrefined ASM generating functions as, respectively,

$$\begin{aligned}
Z_n^{\mathrm{tri}}(x,y;z_1,z_2,z_3)&=Z_n^{\mathrm{quad}}(x,y;z_1,1,z_2,z_3)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{\rho_{\mathrm{B}}(A)}\,z_3^{\rho_{\mathrm{L}}(A)},\\
Z_n^{\mathrm{adj}}(x,y;z_1,z_2)&=Z_n^{\mathrm{quad}}(x,y;z_1,1,1,z_2)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{\rho_{\mathrm{L}}(A)},\\
Z_n^{\mathrm{opp}}(x,y;z_1,z_2)&=Z_n^{\mathrm{quad}}(x,y;z_1,1,z_2,1)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{\rho_{\mathrm{B}}(A)},\\
Z_n(x,y;z)&=Z_n^{\mathrm{quad}}(x,y;z,1,1,1)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z^{\rho_{\mathrm{T}}(A)},\\
Z_n(x,y)&=Z_n^{\mathrm{quad}}(x,y;1,1,1,1)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)},
\end{aligned} \tag{9}$$

where z is a further indeterminate.

Also define alternative quadruply-refined and alternative adjacent-boundary doubly-refined ASM generating functions as, respectively,

$$\begin{aligned}
\widetilde{Z}_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)&=(z_2z_4)^{n-1}\,Z_n^{\mathrm{quad}}(x,y;z_1,\tfrac{1}{z_2},z_3,\tfrac{1}{z_4})\\
&=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{n-\rho_{\mathrm{R}}(A)-1}\,z_3^{\rho_{\mathrm{B}}(A)}\,z_4^{n-\rho_{\mathrm{L}}(A)-1},\\
\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)&=Z_n^{\mathrm{quad}}(x,y;z_1,z_2,1,1)=\textstyle\sum_{A\in\mathrm{ASM}(n)}x^{\nu(A)}\,y^{\mu(A)}\,z_1^{\rho_{\mathrm{T}}(A)}\,z_2^{\rho_{\mathrm{R}}(A)}.
\end{aligned} \tag{10}$$

Note that $\widetilde{Z}_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ is a generating function in which the positions of the 1's in the first and last columns of an ASM are measured relative to the opposite ends of the columns to those used in (4)–(5), i.e., in this generating function, the statistics associated with z_2 and z_4 are, respectively, the numbers of 0's above the 1 in the right-most column, and below the 1 in the left-most column of an ASM.

It can be seen immediately that some relations among generating functions of (9)–(10), involving specializations of boundary parameters to 1, are

$$\begin{aligned}
Z_n^{\text{tri}}(x,y;z_1,1,z_2) &= Z_n^{\text{adj}}(x,y;z_1,z_2),\\
Z_n^{\text{tri}}(x,y;1,z_1,z_2) &= \widetilde{Z}_n^{\text{adj}}(x,y;z_1,z_2),\\
Z_n^{\text{tri}}(x,y;z_1,z_2,1) &= Z_n^{\text{opp}}(x,y;z_1,z_2),\\
Z_n^{\text{adj}}(x,y;z,1) = \widetilde{Z}_n^{\text{adj}}(x,y;z,1) &= Z_n^{\text{opp}}(x,y;z,1) = Z_n(x,y;z),\\
Z_n(x,y;1) &= Z_n(x,y).
\end{aligned}\tag{11}$$

Some elementary identities satisfied by the ASM generating functions of (7), (9) and (10), which follow from the ASM properties outlined in Section 2.1, will now be obtained.

Note that, in this and all subsequent sections, many of the identities which contain the positive integer n will be valid only for all $n \geq 2$, or for all $n \geq 3$. This will often be due to their containing terms (e.g., $Z_{n-1}(x,y)$ or $Z_{n-2}(x,y)$) which are not defined if n is taken to be 1 or 2.

Note also that many of the identities of this section, obtained here using simple combinatorial arguments, can alternatively be obtained as special cases of more general identities which will be derived in subsequent sections using other methods.

Some symmetry relations, which can be derived by acting on ASM(n) with transposition or anticlockwise quarter-turn rotation, and using (6), are

$$\begin{aligned}
Z_n^{\text{quad}}(x,y;z_1,z_2,z_3,z_4) &= Z_n^{\text{quad}}(x,y;z_4,z_3,z_2,z_1)\\
&= x^{n(n-1)/2}\,(z_1z_2z_3z_4)^{n-1}\,Z_n^{\text{quad}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_2},\tfrac{1}{z_3},\tfrac{1}{z_4},\tfrac{1}{z_1}\big),\\
\widetilde{Z}_n^{\text{quad}}(x,y;z_1,z_2,z_3,z_4) &= (z_1z_2z_3z_4)^{n-1}\,\widetilde{Z}_n^{\text{quad}}(x,y;\tfrac{1}{z_4},\tfrac{1}{z_3},\tfrac{1}{z_2},\tfrac{1}{z_1})\\
&= x^{n(n-1)/2}\,\widetilde{Z}_n^{\text{quad}}\big(\tfrac{1}{x},\tfrac{y}{x};z_2,z_3,z_4,z_1\big),\\
Z_n^{\text{tri}}(x,y;z_1,z_2,z_3) &= x^{n(n-1)/2}\,(z_1z_2z_3)^{n-1}\,Z_n^{\text{tri}}(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_2},\tfrac{1}{z_1},\tfrac{1}{z_3}),\\
Z_n^{\text{adj}}(x,y;z_1,z_2) &= Z_n^{\text{adj}}(x,y;z_2,z_1)\\
&= x^{n(n-1)/2}\,(z_1z_2)^{n-1}\,\widetilde{Z}_n^{\text{adj}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_1},\tfrac{1}{z_2}\big),\\
Z_n^{\text{opp}}(x,y;z_1,z_2) &= Z_n^{\text{opp}}(x,y;z_2,z_1)\\
&= x^{n(n-1)/2}\,(z_1z_2)^{n-1}\,Z_n^{\text{opp}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_1},\tfrac{1}{z_2}\big),\\
Z_n(x,y;z) &= x^{n(n-1)/2}\,z^{n-1}\,Z_n\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z}\big),\\
Z_n(x,y) &= x^{n(n-1)/2}\,z^{n-1}\,Z_n\big(\tfrac{1}{x},\tfrac{y}{x}\big).
\end{aligned}\tag{12}$$

Some identities involving specializations of boundary parameters to 0, which follow from the properties of ASMs with a 1 in a corner, as discussed near the end of Section 2.1, are

$$\begin{gathered}
Z_n^{\text{quad}}(x,y;z_1,0,z_2,z_3) = Z_n^{\text{tri}}(x,y;z_1,0,z_3) = Z_{n-1}^{\text{adj}}(x,y;z_1,z_3),\\
\widetilde{Z}_n^{\text{adj}}(x,y;z,0) = Z_n^{\text{opp}}(x,y;z,0) = Z_{n-1}(x,y;z),\\
Z_n^{\text{adj}}(x,y;z,0) = Z_n(x,y;0) = Z_{n-1}(x,y). \qquad (13)
\end{gathered}$$

It will sometimes be useful to refer to boundary parameter coefficients in the adjacent-boundary doubly-refined and singly-refined ASM generating functions. In particular, define

$$\begin{gathered}
Z_n^{\text{adj}}(x,y)_{k_1,k_2} = \text{coefficient of } z_1^{k_1} z_2^{k_2} \text{ in } Z_n^{\text{adj}}(x,y;z_1,z_2),\\
Z_n(x,y)_k = \text{coefficient of } z^k \text{ in } Z_n(x,y;z). \qquad (14)
\end{gathered}$$

These coefficients satisfy identities which correspond to identities of (11)–(13). For example,

$$\begin{gathered}
\textstyle\sum_{k_2=0}^{n-1} Z_n^{\text{adj}}(x,y)_{k_1,k_2} = Z_n(x,y)_{k_1}, \qquad \sum_{k=0}^{n-1} Z_n(x,y)_k = Z_n(x,y),\\
Z_n^{\text{adj}}(x,y)_{k_1,k_2} = Z_n^{\text{adj}}(x,y)_{k_2,k_1}, \qquad Z_n(x,y)_k = x^{n(n-1)/2}\, Z_n(\tfrac{1}{x},\tfrac{y}{x})_{n-1-k},\\
Z_n^{\text{adj}}(x,y)_{k,0} = Z_{n-1}(x,y)\,\delta_{k,0}, \qquad Z_n^{\text{adj}}(x,y)_{k+1,n-1} = x^{n-1}\, Z_{n-1}(x,y)_k,\\
Z_n(x,y)_0 = Z_{n-1}(x,y). \qquad (15)
\end{gathered}$$

Furthermore, the properties of ASMs in which a 1 on a boundary is separated from a corner by a single zero, as discussed at the end of Section 2.1, lead to identities such as

$$\begin{gathered}
Z_n^{\text{adj}}(x,y)_{k,1} = xZ_{n-1}(x,y)_{k-1} + y\,\textstyle\sum_{i=k}^{n-2} Z_{n-1}(x,y)_i - yZ_{n-1}(x,y)\delta_{k,0},\\
Z_n(x,y)_1 = xZ_{n-1}(x,y) + y\,\textstyle\sum_{k=1}^{n-2} k\, Z_{n-1}(x,y)_k. \qquad (16)
\end{gathered}$$

3. Previously-known and Related Results

In this section, an account is given of results for the exact enumeration of $n \times n$ ASMs (for arbitrary, finite n), involving any of the six statistics of (3)–(4), or any of the associated generating functions of (7), (9) or (10). Each of these results is either previously-known, or can be obtained straightforwardly from previously-known results.

Various cases for the values of the parameters x and y, associated with the bulk statistics (3), are first considered separately: $y = 0$ in Section 3.1, $y = x + 1$ in Section 3.2, $x = y = 1$ in Section 3.3, and x and y arbitrary in Section 3.4. Further results are then discussed in Sections 3.5-3.15.

3.1. Bulk parameter $y = 0$.

The case in which the bulk parameter y is 0 corresponds simply to the enumeration of permutation matrices, with prescribed values of the inversion number of associated permutations, and prescribed positions of 1's on the boundaries of the matrices.

Let x-numbers and the x-factorial be defined, as usual, as $[n]_x = 1 + x + \ldots + x^{n-1}$, and $[n]_x! = [n]_x[n-1]_x \ldots [1]_x$.

For $y=0$, the quadruply-refined ASM generating function is explicitly

$$\begin{aligned}
&Z_n^{\mathrm{quad}}(x,0;z_1,z_2,z_3,z_4)=\\
&\quad x^2z_1z_2z_3z_4\textstyle\sum_{0\le i<j\le n-3}\bigl(x^{n+i-j-3}z_4^iz_2^{n-j-3}+x^{n-i+j-4}z_4^{n-i-3}z_2^j\bigr)\times\\
&\qquad\qquad\textstyle\sum_{0\le i<j\le n-3}\bigl(x^{n+i-j-3}z_1^iz_3^{n-j-3}+x^{n-i+j-4}z_1^{n-i-3}z_3^j\bigr)\,[n-4]_x!+\\
&\quad\bigl(xz_4z_1\,[n-2]_{xz_4}\,[n-2]_{xz_1}+z_1z_2(xz_3z_4)^{n-1}\,[n-2]_{xz_1}\,[n-2]_{xz_2}+\\
&\qquad xz_2z_3\,[n-2]_{xz_2}\,[n-2]_{xz_3}+z_3z_4(xz_1z_2)^{n-1}\,[n-2]_{xz_3}\,[n-2]_{xz_4}\bigr)\,[n-3]_x!+\\
&\qquad\qquad\bigl(1+x^{2n-3}(z_1z_2z_3z_4)^{n-1}\bigr)\,[n-2]_x!,
\end{aligned}\tag{17}$$

and the ASM generating functions of (9) are explicitly

$$\begin{aligned}
Z_n^{\mathrm{tri}}(x,0;z_1,z_2,z_3)&=\bigl(xz_2\textstyle\sum_{0\le i<j\le n-3}\bigl(x^{n+i-j-3}z_1^iz_2^{n-j-3}+x^{n-i+j-4}z_1^{n-i-3}z_2^j\bigr)+\\
&\qquad[n-2]_{xz_1}+z_2(xz_1)^{n-2}\,[n-2]_{xz_2}\bigr)\,xz_1z_3\,[n-2]_{xz_3}\,[n-3]_x!+\\
&\quad\bigl(z_1(xz_2z_3)^{n-1}\,[n-2]_{xz_1}+xz_2\,[n-2]_{xz_2}+1+x^{2n-3}(z_1z_2z_3)^{n-1}\bigr)\,[n-2]_x!\\
Z_n^{\mathrm{adj}}(x,0;z_1,z_2)&=xz_1z_2\,[n-1]_{xz_1}\,[n-1]_{xz_2}\,[n-2]_x!+[n-1]_x!,\\
Z_n^{\mathrm{opp}}(x,0;z_1,z_2)&=\textstyle\sum_{0\le i<j\le n-1}\bigl(x^{n+i-j-1}\,z_1^i\,z_2^{n-j-1}+x^{n-i+j-2}\,z_1^{n-i-1}\,z_2^j\bigr)\,[n-2]_x!,\\
Z_n(x,0;z)&=[n]_{xz}\,[n-1]_x!,\\
Z_n(x,0)&=[n]_x!.
\end{aligned}\tag{18}$$

Each of these formulae is either a standard result for permutations, or a straightforward variation of such a result, and each formula can be derived using simple combinatorial arguments. See, for example, Stanley [132, Cor. 1.3.13] for a derivation of the last equation of (18).

Note that if (17) is shown first to be valid, then each equation of (18) can subsequently be obtained by setting certain boundary parameters in (17) to 1. Alternatively, if the last two equations of (18) are shown first to be valid, then (17) and the remaining equations of (18) can subsequently be obtained by using the general results (79), (81) (or (84)), (85) and (86), which will be given in Section 4.

In interpreting (17), note also that the seven main terms on the RHS of (17) (i.e., the term ending in $[n-4]_x!$, the four terms ending in $[n-3]_x!$ and the two terms ending in $[n-2]_x!$) correspond to sums over the sets of $n\times n$ permutation matrices A with $(A_{11},A_{1n},A_{nn},A_{n1})$ equal to $(0,0,0,0)$, $(0,0,1,0)$, $(0,0,0,1)$, $(1,0,0,0)$, $(0,1,0,0)$, $(1,0,1,0)$ or $(0,1,0,1)$, respectively. In subsequent formulae for quadruply-refined ASM generating functions (e.g., (19), (48) and (79)), the RHS will again consist of seven or eight main terms, but these terms no longer correspond simply to sums over sets of ASMs with fixed values of the four corner entries.

3.2. **Bulk parameters satisfying $y=x+1$.** The case in which the bulk parameters are related by $y=x+1$ is closely related to x-determinants (or, in the notation usually used, λ-determinants) of matrices, domino tilings of an Aztec diamond, tournaments, and the free fermion case of the six-vertex model. (Note that the sub-case $x = 1$ and $y = 2$, which corresponds to the so-called 2-enumeration of ASMs, is often considered separately.) Due to certain combinatorial and algebraic simplifications, it is again possible to obtain explicit formulae.

For $y = x + 1$, the quadruply-refined ASM generating function is given explicitly by

$$\begin{aligned}
&(xz_4z_1+z_4+z_1-1)(xz_1z_2-xz_1-xz_2-1)\times\\
&\qquad(xz_2z_3+z_2+z_3-1)(xz_3z_4-xz_3-xz_4-1)\,Z_n^{\mathrm{quad}}(x,x+1;z_1,z_2,z_3,z_4)=\\
&z_1z_2z_3z_4(xz_1z_3+1)(xz_2z_4+1)\big((xz_1+1)(xz_2+1)(xz_3+1)(xz_4+1)\big)^{n-2}(x+1)^{(n-4)(n-5)/2}-\\
&\qquad\Big((z_2-1)(z_3-1)z_4z_1\big((xz_4+1)(xz_1+1)\big)^{n-2}+\\
&\qquad\qquad(z_4-1)(z_1-1)z_2z_3\big((xz_2+1)(xz_3+1)\big)^{n-2}\Big)\times\\
&\qquad\qquad(xz_1z_2-xz_1-xz_2-1)(xz_3z_4-xz_3-xz_4-1)(x+1)^{(n-3)(n-4)/2}-\\
&\qquad\Big((z_3-1)(z_4-1)z_1z_2(z_3z_4)^{n-1}\big((xz_1+1)(xz_2+1)\big)^{n-2}+\\
&\qquad\qquad(z_1-1)(z_2-1)z_3z_4(z_1z_2)^{n-1}\big((xz_3+1)(xz_4+1)\big)^{n-2}\Big)\times\\
&\qquad\qquad(xz_4z_1+z_4+z_1-1)(xz_2z_3+z_2+z_3-1)\,x^n(x+1)^{(n-3)(n-4)/2}+\\
&\quad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\Big((xz_1z_2-xz_1-xz_2-1)(xz_3z_4-xz_3-xz_4-1)+\\
&\qquad(xz_4z_1+z_4+z_1-1)(xz_2z_3+z_2+z_3-1)(z_1z_2z_3z_4)^{n-1}x^{2n-1}\Big)(x+1)^{(n-2)(n-3)/2},
\end{aligned}\tag{19}$$

and the ASM generating functions of (9) are given explicitly by

$$\begin{aligned}
&(xz_1z_3+z_1+z_3-1)(xz_2z_3-xz_2-xz_3-1)Z_n^{\mathrm{tri}}(x,x+1;z_1,z_2,z_3)=\\
&\quad-z_1z_3(xz_1z_2+1)\big((xz_1+1)(xz_2+1)\big)^{n-2}(xz_3+1)^{n-1}(x+1)^{(n-3)(n-4)/2}+\\
&\quad(z_2-1)(z_3-1)(xz_1z_3+z_1+z_3-1)z_1(z_2z_3)^{n-1}(xz_1+1)^{n-2}x^n(x+1)^{(n-2)(n-3)/2}-\\
&\quad(z_1-1)(z_3-1)(xz_2z_3-xz_2-xz_3-1)(xz_2+1)^{n-2}(x+1)^{(n-2)(n-3)/2},\\
&(xz_1z_2+z_1+z_2-1)Z_n^{\mathrm{adj}}(x,x+1;z_1,z_2)=z_1z_2\big((xz_1+1)(xz_2+1)\big)^{n-1}(x+1)^{(n-2)(n-3)/2}\\
&\qquad\qquad-(z_1-1)(z_2-1)\,(x+1)^{(n-1)(n-2)/2},\\
&Z_n^{\mathrm{opp}}(x,x+1;z_1,z_2)=(xz_1z_2+1)\,\big((xz_1+1)(xz_2+1)\big)^{n-2}\,(x+1)^{(n-2)(n-3)/2},\\
&Z_n(x,x+1;z)=(xz+1)^{n-1}\,(x+1)^{(n-1)(n-2)/2},\\
&Z_n(x,x+1)=(x+1)^{n(n-1)/2}.
\end{aligned}\tag{20}$$

Each of these formulae is either a previously-known result, or a straightforward variation of such a result. A derivation of these formulae, based on a result of Robbins and Rumsey [125, Sec. 5] for x-determinants, will now be given, since this derivation involves the use of a quadratic relation satisfied by the quadruply-refined ASM generating function, and thereby shares some features with new material which will be presented in Section 4. An alternative derivation, based on the Izergin–Korepin formula for the partition function of six-vertex model with DWBC and the Cauchy double alternant evaluation, will be given in Section 5.8.

For an $n \times n$ matrix of indeterminates, $\bigl(M_{ij}\bigr)_{1\le i,j\le n}$, define

$$\mathcal{Z}_n(x,M) = \textstyle\sum_{A\in\mathrm{ASM}(n)} x^{\nu(A)}\,(x+1)^{\mu(A)}\,\prod_{i,j=1}^{n}(M_{ij})^{A_{ij}}, \tag{21}$$

this being the so-called x-determinant of M, as introduced by Robbins and Rumsey [125, Sec. 5]. It can be seen that $\mathcal{Z}_n(-1,M)$ is the standard determinant of M, since the RHS of (21) is then a sum over all $n\times n$ permutation matrices A, with $\nu(A)$ being the number of inversions in the permutation associated with A.

It follows from a result of Robbins and Rumsey [125, Sec. 5] that

$$\mathcal{Z}_n(x,M)\,\mathcal{Z}_{n-2}(x,M_{\mathrm{C}}) = \mathcal{Z}_{n-1}(x,M_{\mathrm{TL}})\,\mathcal{Z}_{n-1}(x,M_{\mathrm{BR}})+x\,\mathcal{Z}_{n-1}(x,M_{\mathrm{TR}})\,\mathcal{Z}_{n-1}(x,M_{\mathrm{BL}}), \tag{22}$$

where M_{TL}, M_{TR}, M_{BR} and M_{BL} denote the $(n-1)\times(n-1)$ submatrices corresponding to the top-left, top-right, bottom-right and bottom-left corners of M, respectively, and M_{C} denotes the central $(n-2)\times(n-2)$ submatrix of M. For $x=-1$, (22) is the Desnanot-Jacobi determinant identity, which will be discussed in more detail in Section 5.5.

Taking M in (22) to be

$$M = \begin{pmatrix} 1 & z_1 & \dots & z_1^{n-2} & (z_1z_2)^{n-1} \\ z_4 & 1 & \dots & 1 & z_2^{n-2} \\ \vdots & \vdots & & \vdots & \vdots \\ z_4^{n-2} & 1 & \dots & 1 & z_2 \\ (z_3z_4)^{n-1} & z_3^{n-2} & \dots & z_3 & 1 \end{pmatrix}, \tag{23}$$

it follows, using definitions from (7), (9), (10) and (21), that the quadruply-refined ASM generating function satisfies the quadratic relation

$$\begin{aligned} Z_n^{\mathrm{quad}}(x,x{+}1;z_1,z_2,z_3,z_4)\,Z_{n-2}(x,x{+}1) = Z_{n-1}^{\mathrm{adj}}(x,x{+}1;z_4,z_1)\,Z_{n-1}^{\mathrm{adj}}(x,x{+}1;z_2,z_3) + {} \\ xz_1z_2z_3z_4\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,x{+}1;z_1,z_2)\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,x{+}1;z_3,z_4). \end{aligned} \tag{24}$$

Note that a quadratic relation for arbitrary x and y will be given in (79), and that a different quadratic relation for the present case can be obtained by setting $y=x+1$ in (79).

Setting $z_1=z_2=z_3=z_4=1$ in (24) gives

$$Z_n(x,x{+}1)\,Z_{n-2}(x,x{+}1) = (x{+}1)\,Z_{n-1}(x,x{+}1)^2, \tag{25}$$

which, together with $Z_1(x,x{+}1)=1$ and $Z_2(x,x{+}1)=x{+}1$, gives the formula in (20) for $Z_n(x,x{+}1)$.

Setting $z_2=z_3=z_4=1$ in (24), and relabeling z_1 as z, gives

$$Z_n(x,x{+}1;z)\,Z_{n-2}(x,x{+}1) = (xz{+}1)\,Z_{n-1}(x,x{+}1;z)\,Z_{n-1}(x,x{+}1), \tag{26}$$

which, together with $Z_1(x,x{+}1;z)=1$ and the formula for $Z_n(x,x{+}1)$, gives the formula in (20) for $Z_n(x,x{+}1;z)$. The formula for $Z_n^{\mathrm{opp}}(x,x{+}1;z_1,z_2)$ can be obtained similarly, by setting $z_2=z_4=1$ in (24), and using the formulae for $Z_n(x,x{+}1;z)$ and $Z_n(x,x{+}1)$.

Setting $z_2=z_3=1$ in (24), and relabeling z_4 as z_2, gives

$$\begin{aligned}&Z^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)\,Z_{n-2}(x,x{+}1)=\\&\qquad Z^{\mathrm{adj}}_{n-1}(x,x{+}1;z_1,z_2)\,Z_{n-1}(x,x{+}1)+xz_1z_2\,Z_{n-1}(x,x{+}1;z_1)\,Z_{n-1}(x,x{+}1;z_2),\end{aligned}\tag{27}$$

which, together with $Z^{\mathrm{adj}}_1(x,x{+}1;z_1,z_2)=1$, gives

$$Z^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)=Z_{n-1}(x,x{+}1)\left(1+xz_1z_2\sum_{i=1}^{n-1}\frac{Z_i(x,x{+}1;z_1)\,Z_i(x,x{+}1;z_2)}{Z_{i-1}(x,x{+}1)\,Z_i(x,x{+}1)}\right),\tag{28}$$

where, in the sum over i, $Z_0(x,x{+}1)$ is taken to be 1. Substituting the formulae for $Z_i(x,x{+}1;z)$ and $Z_i(x,x{+}1)$ into (28), and then performing the sum over i and simplifying, gives the formula in (20) for $Z^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)$.

Finally, the formulae in (19)–(20) for $Z^{\mathrm{quad}}_n(x,x{+}1;z_1,z_2,z_3,z_4)$ and $Z^{\mathrm{tri}}_n(x,x{+}1;z_1,z_2,z_3)$ can be obtained using (24), together with the formulae for $Z^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)$ and $Z_n(x,x{+}1)$, and a similar formula for $\widetilde{Z}^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)$ (which can be obtained from the formula for $Z^{\mathrm{adj}}_n(x,x{+}1;z_1,z_2)$ and the relevant equation from (12)).

For further information regarding various aspects of the case $y=x+1$ and closely related cases, including details of several alternative methods for obtaining formulae such as (19)–(20), see, for example, Bogoliubov, Pronko and Zvonarev [20, Sec. 5], Bousquet-Mélou and Habseiger [21, Sec. 5], Bressoud [23], Brualdi and Kirkland [26], Brubaker, Bump and Friedberg [28, 29], Bump, McNamara and Nakasuji [30], Chapman [33], Ciucu [36], [37, Thm. 6.1], Colomo and Pronko [45, Secs. 4.2 & 5.2], [46, Sec. 4.3], Elkies, Kuperberg, Larsen and Propp [63, 64], Eu and Fu [65], Ferrari and Spohn [66], Hamel and King [77, Cor. 5.1], Kuo [91, Sec. 3], Kuperberg [93, Thm. 3], Lascoux [97], McNamara [101], Mills, Robbins and Rumsey [103, Sec. 6], Okada [111, Thm. 2.4(1), third eq.], Rosengren [126, Sec. 9], Striker [134, Sec. 5], [136, Sec. 6], and Tokuyama [142, Cor. 3.4].

3.3. **Bulk parameters $x=y=1$.**

The case in which the bulk parameters x and y are both 1 involves numbers of ASMs with prescribed configurations on certain boundaries, but without prescribed values of bulk statistics. The cases of zero, one or two boundaries will be considered together first, followed by the cases of three or four boundaries.

Define unrefined, singly-refined, opposite-boundary doubly-refined and adjacent-boundary doubly-refined ASM numbers as, respectively,

$$\begin{aligned}\mathcal{A}_n&=|\mathrm{ASM}(n)|,\\\mathcal{A}_{n,k}&=|\{A\in\mathrm{ASM}(n)\mid A_{1,k+1}=1\}|,\\\mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2}&=|\{A\in\mathrm{ASM}(n)\mid A_{1,k_1+1}=A_{n,n-k_2}=1\}|,\\\mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2}&=|\{A\in\mathrm{ASM}(n)\mid A_{1,k_1+1}=A_{k_2+1,1}=1\}|,\end{aligned}\tag{29}$$

for $0 \leq k, k_1, k_2 \leq n-1$, with the numbers being 0 for k, k_1 or k_2 outside this range. These numbers are therefore related to generating functions of (9)–(10) by

$$\begin{aligned} Z_n(1,1) &= \mathcal{A}_n, \\ Z_n(1,1;z) &= \textstyle\sum_{k=0}^{n-1} \mathcal{A}_{n,k}\, z^k, \\ Z_n^{\mathrm{opp}}(1,1;z_1,z_2) &= \textstyle\sum_{k_1,k_2=0}^{n-1} \mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2}\, z_1^{k_1}\, z_2^{k_2}, \\ Z_n^{\mathrm{adj}}(1,1;z_1,z_2) &= \textstyle\sum_{k_1,k_2=0}^{n-1} \mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2}\, z_1^{k_1}\, z_2^{k_2}, \\ \widetilde{Z}_n^{\mathrm{adj}}(1,1;z_1,z_2) &= \textstyle\sum_{k_1,k_2=0}^{n-1} \mathcal{A}^{\mathrm{adj}}_{n,n-1-k_1,n-1-k_2}\, z_1^{k_1}\, z_2^{k_2}, \end{aligned} \tag{30}$$

and to the boundary-parameter coefficients of (14) by

$$\mathcal{A}_{n,k} = Z_n(1,1)_k, \qquad \mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2} = Z_n^{\mathrm{adj}}(1,1)_{k_1,k_2}. \tag{31}$$

The ASM numbers of (29) satisfy various elementary identities related to their definitions, symmetry properties of ASMs or properties of ASMs with a 1 in a corner, and which correspond to identities, such as (11)–(13) and (15), satisfied by the generating functions. Some examples of these identities are

$$\begin{gathered} \textstyle\sum_{k=0}^{n-1} \mathcal{A}_{n,k} = \mathcal{A}_n, \qquad \mathcal{A}_{n,k} = \mathcal{A}_{n,n-1-k}, \qquad \mathcal{A}_{n,0} = \mathcal{A}_{n-1}, \\ \textstyle\sum_{k_2=0}^{n-1} \mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2} = \sum_{k_2=0}^{n-1} \mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2} = \mathcal{A}_{n,k_1}, \\ \mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2} = \mathcal{A}^{\mathrm{opp}}_{n,k_2,k_1} = \mathcal{A}^{\mathrm{opp}}_{n,n-1-k_1,n-1-k_2}, \qquad \mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2} = \mathcal{A}^{\mathrm{adj}}_{n,k_2,k_1}, \\ \mathcal{A}^{\mathrm{adj}}_{n,k,0} = \mathcal{A}_{n-1}\, \delta_{k,0}, \qquad \mathcal{A}^{\mathrm{opp}}_{n,k,0} = \mathcal{A}^{\mathrm{adj}}_{n,k+1,n-1} = \mathcal{A}_{n-1,k}. \end{gathered} \tag{32}$$

Furthermore, some identities related to properties of ASMs in which a 1 on a boundary is separated from a corner by a single zero, and which can be obtained from (16) and the first two identities of (32), are

$$\mathcal{A}^{\mathrm{adj}}_{n,k,1} = \textstyle\sum_{i=k-1}^{n-2} \mathcal{A}_{n-1,i} - \mathcal{A}_{n-1}\, \delta_{k,0}, \qquad \mathcal{A}_{n,1} = \mathcal{A}_{n-1} + \sum_{k=1}^{n-2} k\, \mathcal{A}_{n-1,k} = \tfrac{n}{2}\, \mathcal{A}_{n-1}.$$

Proceeding to more general identities, for which derivations using combinatorial arguments are not currently known, formulae which give the ASM numbers of (29) explicitly are

$$\mathcal{A}_n = \textstyle\prod_{i=0}^{n-1} \frac{(3i+1)!}{(n+i)!}, \tag{33}$$

$$\mathcal{A}_{n,k} = \begin{cases} \frac{(n+k-1)!\,(2n-k-2)!}{k!\,(n-k-1)!\,(2n-2)!} \prod_{i=0}^{n-2} \frac{(3i+1)!}{(n+i-1)!}, & 0 \leq k \leq n-1 \\ 0, & \text{otherwise}, \end{cases} \tag{34}$$

$$\begin{aligned} \mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2} = \tfrac{1}{\mathcal{A}_{n-1}} \textstyle\sum_{i=0}^{\min(k_1,n-k_2-1)} \big(&\mathcal{A}_{n,k_1-i}\, \mathcal{A}_{n-1,k_2+i} + \mathcal{A}_{n-1,k_1-i-1}\, \mathcal{A}_{n,k_2+i} - {} \\ &\mathcal{A}_{n,k_1-i-1}\, \mathcal{A}_{n-1,k_2+i} - \mathcal{A}_{n-1,k_1-i-1}\, \mathcal{A}_{n,k_2+i+1}\big), \end{aligned} \tag{35}$$

$$\mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2} = \begin{cases} \mathcal{A}_{n-1}, & k_1 = k_2 = 0 \\ \binom{k_1+k_2-2}{k_1-1} \mathcal{A}_{n-1} - \sum_{i=1}^{k_1} \sum_{j=1}^{k_2} \binom{k_1+k_2-i-j}{k_1-i} \mathcal{A}^{\mathrm{opp}}_{n,i-1,n-j}, & 1 \leq k_1, k_2 \leq n-1 \\ 0, & \text{otherwise}, \end{cases} \tag{36}$$

It follows from (33)–(34) that the unrefined and singly-refined ASM numbers satisfy simple recursion relations, such as

$$\begin{aligned}\mathcal{A}_n &= \tfrac{(n-1)!\,(3n-2)!}{(2n-2)!\,(2n-1)!}\,\mathcal{A}_{n-1},\\ \mathcal{A}_n\,\mathcal{A}_{n-2} &= \tfrac{3(3n-2)(3n-4)}{4(2n-1)(2n-3)}\,\mathcal{A}_{n-1}^2,\\ k(2n-k-1)\mathcal{A}_{n,k} &= (n-k)(n+k-1)\,\mathcal{A}_{n,k-1},\end{aligned}\tag{37}$$

and that the singly-refined ASM generating function at $x=y=1$ can be expressed in terms of the Gaussian hypergeometric function as

$$Z_n(1,1;z)=\mathcal{A}_{n-1}\,{}_2F_1\!\left[\begin{matrix}1-n,\ n\\ 2-2n\end{matrix};\,z\right],\tag{38}$$

and satisfies the hypergeometric differential equation

$$z(1-z)\tfrac{d^2}{dz^2}Z_n(1,1;z)+2(1-n-z)\tfrac{d}{dz}Z_n(1,1;z)+n(n-1)Z_n(1,1;z)=0.\tag{39}$$

The singly-refined ASM numbers also satisfy, for $0\le k\le n-1$,

$$\mathcal{A}_{n,k}=\textstyle\sum_{i=0}^{k}(-1)^i\binom{n+k-1}{k-i}\mathcal{A}_{n,i}.\tag{40}$$

Recursion relations which involve the opposite-boundary and adjacent-boundary doubly-refined ASM numbers are

$$\begin{aligned}(\mathcal{A}^{\mathrm{opp}}_{n,k_1-1,k_2}-\mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2-1})\,\mathcal{A}_{n-1} &= \mathcal{A}_{n,k_1-1}\,\mathcal{A}_{n-1,k_2-1}-\mathcal{A}_{n,k_1}\,\mathcal{A}_{n-1,k_2-1}-{}\\ &\qquad \mathcal{A}_{n-1,k_1-1}\,\mathcal{A}_{n,k_2-1}+\mathcal{A}_{n-1,k_1-1}\,\mathcal{A}_{n,k_2},\end{aligned}\tag{41}$$

$$\mathcal{A}^{\mathrm{adj}}_{n,k_1-1,k_2}+\mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2-1}-\mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2}=\mathcal{A}^{\mathrm{opp}}_{n,k_1-1,n-k_2}-(\delta_{k_1,1}-\delta_{k_1,0})(\delta_{k_2,1}-\delta_{k_2,0})\,\mathcal{A}_{n-1}.\tag{42}$$

It follows that, in terms of generating functions, (41) can be written as

$$\begin{aligned}(z_1-z_2)\,Z^{\mathrm{opp}}_n(1,1;z_1,z_2)\,\mathcal{A}_{n-1} &= (z_1-1)\,z_2\,Z_n(1,1;z_1)\,Z_{n-1}(1,1;z_2)-{}\\ &\qquad z_1\,(z_2-1)\,Z_{n-1}(1,1;z_1)\,Z_n(1,1;z_2),\end{aligned}\tag{43}$$

and (42) can be written as

$$(z_1+z_2-1)\,Z^{\mathrm{adj}}_n(1,1;z_1,z_2)=z_1\,z_2^n\,Z^{\mathrm{opp}}_n(1,1;z_1,\tfrac{1}{z_2})-(z_1-1)(z_2-1)\,\mathcal{A}_{n-1}.\tag{44}$$

The relation (44) will be used in Section 4.3 for the derivation of results for the quadruply-refined ASM generating function at $x=y=1$.

The opposite-boundary doubly-refined ASM generating function at $x=y=1$ can also be expressed as

$$\begin{aligned}Z^{\mathrm{opp}}_n(1,1;z_1,z_2) &= 3^{-n(n-1)/2}\,\big(q^2(z_1+q)(z_2+q)\big)^{n-1}\times\\ &\qquad s_{(n-1,n-1,\ldots,2,2,1,1)}\big(\tfrac{qz_1+1}{z_1+q},\,\tfrac{qz_2+1}{z_2+q},\,\underbrace{1,\ldots,1}_{2n-2}\big)\big|_{q=e^{\pm2\pi i/3}},\end{aligned}\tag{45}$$

where $s_{(n-1,n-1,\ldots,2,2,1,1)}\bigl(\frac{qz_1+1}{z_1+q},\frac{qz_1+1}{z_1+q},1,\ldots,1\bigr)$ is the Schur function indexed by the double-staircase partition $(n-1,n-1,\ldots,2,2,1,1)$, evaluated at the $2n$ parameters $\frac{qz_1+1}{z_1+q}$, $\frac{qz_1+1}{z_1+q}$, $1,\ldots,1$. Setting $z_1=z_2=1$ in (45), and using (30) and (32), it follows that

$$\begin{aligned}\mathcal{A}_n = 3^{-n(n-1)/2}\times\bigl(&\text{number of semistandard Young tableaux of shape}\\ &(n-1,n-1,\ldots,2,2,1,1)\text{ with entries from }\{1,\ldots,2n\}\bigr).\end{aligned}\tag{46}$$

The formula (33) can be obtained from (46) using the hook-content formula for semistandard Young tableaux.

A further identity satisfied by the opposite-boundary doubly-refined and unrefined ASM numbers is

$$\det_{0\le k_1,k_2\le n-1}\bigl(\mathcal{A}^{\rm opp}_{n,k_1,k_2}\bigr)=(-1)^{n(n+1)/2+1}\,(\mathcal{A}_{n-1})^{n-3}.\tag{47}$$

Proceeding now to the enumeration of ASMs with prescribed configurations on three or four boundaries, the alternative quadruply-refined ASM generating function at $x=y=1$ is given by

$$\begin{aligned}&(z_4z_1-z_4+1)(z_1z_2-z_1+1)(z_2z_3-z_2+1)(z_3z_4-z_3+1)\,\widetilde{Z}^{\rm quad}_n(1,1;z_1,z_2,z_3,z_4)=\\
&\quad\frac{z_1\,z_2\,z_3\,z_4\,\det_{1\le i,j\le 4}\bigl(z_i^{j-1}\,(z_i-1)^{4-j}\,Z_{n-j+1}(1,1;z_i)\bigr)}{\mathcal{A}_{n-1}\,\mathcal{A}_{n-2}\,\mathcal{A}_{n-3}\,\prod_{1\le i<j\le 4}(z_i-z_j)}+\\
&\quad(z_2-1)(z_3-1)(z_4z_1-z_4+1)(z_1z_2-z_1+1)(z_3z_4-z_3+1)(z_2z_4)^{n-1}Z^{\rm adj}_{n-1}(1,1;\tfrac{1}{z_4},z_1)+\\
&\quad(z_3-1)(z_4-1)(z_1z_2-z_1+1)(z_2z_3-z_2+1)(z_4z_1-z_4+1)(z_1z_3)^{n-1}Z^{\rm adj}_{n-1}(1,1;\tfrac{1}{z_1},z_2)+\\
&\quad(z_4-1)(z_1-1)(z_2z_3-z_2+1)(z_3z_4-z_3+1)(z_1z_2-z_1+1)(z_2z_4)^{n-1}Z^{\rm adj}_{n-1}(1,1;\tfrac{1}{z_2},z_3)+\\
&\quad(z_1-1)(z_2-1)(z_3z_4-z_3+1)(z_4z_1-z_4+1)(z_2z_3-z_2+1)(z_1z_3)^{n-1}Z^{\rm adj}_{n-1}(1,1;\tfrac{1}{z_3},z_4)-\\
&\qquad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\bigl((z_1z_2-z_1+1)(z_3z_4-z_3+1)(z_2z_4)^{n-1}+\\
&\qquad\qquad(z_2z_3-z_2+1)(z_4z_1-z_4+1)(z_1z_3)^{n-1}\bigr)\mathcal{A}_{n-2}.\end{aligned}\tag{48}$$

Setting $z_2=1$ in (48) (and then relabeling z_3 as z_2 and z_4 as z_3), it follows that the triply-refined ASM generating function at $x=y=1$ is given by

$$\begin{aligned}&(z_1z_3-z_3+1)(z_2z_3-z_2+1)z_3^{n-1}Z^{\rm tri}_n(1,1;z_1,z_2,\tfrac{1}{z_3})=\\
&\quad\frac{z_1\,z_3\,\det_{1\le i,j\le 3}\bigl(z_i^{j-1}\,(z_i-1)^{3-j}\,Z_{n-j+1}(1,1;z_i)\bigr)}{\mathcal{A}_{n-1}\,\mathcal{A}_{n-2}\,\prod_{1\le i<j\le 3}(z_i-z_j)}+\\
&\qquad(z_2-1)(z_3-1)(z_1z_3-z_3+1)z_1z_2^{n-1}Z_{n-1}(1,1;z_1)+\\
&\qquad\qquad(z_1-1)(z_3-1)(z_2z_3-z_2+1)z_3^{n-1}Z_{n-1}(1,1;z_2).\end{aligned}\tag{49}$$

Note that the identities (43) and (44), satisfied by the doubly-refined ASM generating functions at $x=y=1$, can be regarded as special cases of the identity (49) satisfied by the triply-refined ASM generating function. More specifically, setting $z_3=1$ in (49) gives (43), while setting $z_2=1$ in (49), and using (43), gives (44).

Note also that, by using (44) to replace each case of an adjacent-boundary doubly-refined ASM generating function in (48) by an opposite-boundary doubly-refined ASM generating function, (48) can be restated as

$$
\begin{aligned}
&(z_4z_1-z_4+1)(z_1z_2-z_1+1)(z_2z_3-z_2+1)(z_3z_4-z_3+1)\,\widetilde{Z}_n^{\mathrm{quad}}(1,1;z_1,z_2,z_3,z_4) = \\
&\quad \frac{z_1\,z_2\,z_3\,z_4\,\det_{1\le i,j\le 4}\bigl(z_i^{j-1}\,(z_i-1)^{4-j}\,Z_{n-j+1}(1,1;z_i)\bigr)}{\mathcal{A}_{n-1}\,\mathcal{A}_{n-2}\,\mathcal{A}_{n-3}\,\prod_{1\le i<j\le 4}(z_i-z_j)} + \\
&\qquad (z_2-1)(z_3-1)(z_1z_2-z_1+1)(z_3z_4-z_3+1)z_4z_1z_2^{n-1}\,Z_{n-1}^{\mathrm{opp}}(1,1;z_4,z_1) + \\
&\qquad (z_3-1)(z_4-1)(z_2z_3-z_2+1)(z_4z_1-z_4+1)z_1z_2z_3^{n-1}\,Z_{n-1}^{\mathrm{opp}}(1,1;z_1,z_2) + \\
&\qquad (z_4-1)(z_1-1)(z_3z_4-z_3+1)(z_1z_2-z_1+1)z_2z_3z_4^{n-1}\,Z_{n-1}^{\mathrm{opp}}(1,1;z_2,z_3) + \\
&\qquad (z_1-1)(z_2-1)(z_4z_1-z_4+1)(z_2z_3-z_2+1)z_3z_4z_1^{n-1}\,Z_{n-1}^{\mathrm{opp}}(1,1;z_3,z_4) + \\
&\quad (z_1-1)(z_2-1)(z_3-1)(z_4-1)\bigl((z_1z_2-z_1+1)(z_3z_4-z_3+1)(z_2z_4)^{n-1} + \\
&\qquad\qquad (z_2z_3-z_2+1)(z_4z_1-z_4+1)(z_1z_3)^{n-1}\bigr)\mathcal{A}_{n-2}.
\end{aligned}
\tag{50}
$$

The first terms (i.e., the determinant terms) on the RHS of (48)–(50) are proportional to certain cases of a function which will be defined in (69). Specifically, using (69), the first terms on the RHS of (48) (or (50)) and (49) are $z_1z_2z_3z_4\,X_n(1,1;z_1,z_2,z_3,z_4)$ and $z_1z_2z_3\,X_n(1,1;z_1,z_2,z_3)$, respectively. It will also be seen, in (73), that the function (69) at $x=y=1$, and hence the first terms on the RHS of (48)–(50), are related to a certain Schur function.

Further expressions for $\widetilde{Z}_n^{\mathrm{quad}}(1,1;z_1,z_2,z_3,z_4)$ and $Z_n^{\mathrm{tri}}(1,1;z_1,z_2,\frac{1}{z_3})$, which differ from (48)–(49) in the first terms on each RHS, will be obtained in Corollaries 9 and 10.

The origins of the results given in this section will now be outlined.

The product formula (33) was conjectured by Mills, Robbins and Rumsey [102, 103, Conj. 1], and first proved by Zeilberger [145] and, shortly thereafter, but using a different method, by Kuperberg [92]. The product formula (34) was first proved by Zeilberger [146], and confirms the validity of further conjectures of Mills, Robbins and Rumsey [102, 103, Conj. 2].

Alternative proofs of (33)–(34) have been given by Colomo and Pronko [45, Sec. 5.3], [46, Sec. 4.2], Fischer [68], and Stroganov [140, Sec. 4]. (See also Razumov and Stroganov [117, Sec. 2], [121, Sec. 2] for additional details related to the third of these proofs.)

The formulae (33)–(34) also follow from certain other known results. For example, as already indicated, (33) can be obtained from the result (46) and the hook-content formula. Alternatively, (33) can be obtained as a special case of a result of Rosengren [126, Cor. 8.4]. (See also, for example, Rosengren [127] for further related results.) Finally, (34) can be obtained by combining a result of Behrend, Di Francesco and Zinn-Justin [17, Thm. 1] that $\mathcal{A}_{n,k}$ is the number of descending plane partitions with each part at most n and exactly k parts equal to n, with a result of Mills, Robbins and Rumsey [102, Sec. 5] which gives a product formula for the number of such objects.

Among these various derivations of (33)–(34), all make essential use of the Izergin–Korepin formula [81] for the partition function of the six-vertex model with DWBC, or related properties of integrability, except for Zeilberger's proof [145] of (33), which makes essential use of a result of Andrews [4] for the number of totally symmetric self-complementary plane partitions in a $2n \times 2n \times 2n$ box, and Fischer's proof [68], which makes essential use of an operator formula obtained by Fischer [67, 69] (see also Colomo and Pronko [49, Eq. (3.3)]).

The hypergeometric function expression (38) is given by Colomo and Pronko [43, Eq. (2.16)], [45, Eq. (5.43)], [46, Eq. (4.19)], and the differential equation (39) is given by Stroganov [140, Eq. (26)].

The relation (40) was obtained by Fischer [68, Sec. 3], with a corresponding relation for descending plane partitions, containing a further parameter associated with the sum of the parts of a descending plane partition, having been obtained previously by Mills, Robbins and Rumsey [102, Sec. 5]. It was also shown by Fischer [68, Sec. 4] that the formula (34) follows from (40), the first three equations of (32), and $\mathcal{A}_{1,0} = 1$.

The relations (41)–(44) for the opposite-boundary and adjacent-boundary doubly-refined ASM numbers were first obtained by Stroganov [140, Sec. 5].

The formula (35) follows easily from (41). The formula (36) was first obtained by Fischer [71, p. 570], and then Ayyer and Romik [12, p. 4], and can be derived by dividing both sides of (44) by $z_1 + z_2 - 1$ and equating coefficients of $z_1^{k_1} z_2^{k_2}$ on each side.

The Schur function expression (45) was obtained by Di Francesco and Zinn-Justin [59, Eqs. (2.2) & (2.4)], using a result of Okada [111, Thm. 2.4(1), second equation]. A generalization of (45) will be given in (73), and a derivation of (73) will be given in Section 5.10.

The relation (46) was first obtained by Okada [111, Thm. 1.2 (A1)].

The identity (47) was obtained by Biane, Cantini and Sportiello [18, Thm. 1], by combining (45) with a certain Schur function identity, also obtained by Biane, Cantini and Sportiello [18, Thm. 2].

Some further aspects of the opposite-boundary doubly-refined ASM numbers are discussed by Fischer [69, Sec. 6].

Identities for the quadruply- and triply-refined ASM generating functions at $x = y = 1$, which appear to be equivalent to (48)–(49), were obtained recently by Ayyer and Romik [12, Thms. 1–2]. The forms of (48)–(49) given here were observed by Colomo [40], and can be derived by combining a result which will be given in (71), with a result that the partition function of the six-vertex model with DWBC is symmetric in all of its spectral parameters at its so-called combinatorial point. A derivation of (48) using this approach will be given in Section 5.10. (Roughly speaking, this derivation involves associating spectral parameters t_1, t_2, t_3 and t_4 with the first row, last column, last row and first column, respectively, of an ASM. The symmetry of the partition function in all of its spectral parameters then enables t_2 and t_4 to be associated instead with rows, so that the result (71), which involves parameters associated with several rows, can be applied.)

Note that, since setting certain boundary parameters in (48) to 1 gives (43), (44) and (49), Section 5.10 also provides derivations of these other identities. (For the identity (44), such a derivation essentially depends only the symmetry of the partition function in all of its spectral parameters, and not on (71).)

In the versions of (48)–(49) obtained by Ayyer and Romik [12, Thms. 1–2], slightly different quadruply-refined and triply-refined ASM generating functions are used, and matrices are given which seem to be related by column operations to those in the first terms on the RHS of (48)–(49). In particular, the quadruply-refined generating function used there is $Z'_n(x,y;z_1,z_2,z_3,z_4) = \sum_{\substack{A\in\mathrm{ASM}(n)\\ A_{11}=A_{1n}=A_{nn}=A_{n1}=0}} x^{\nu(A)}\, y^{\mu(A)}\, z_1^{\rho_{\mathrm{T}}(A)-1}\, z_2^{n-2-\rho_{\mathrm{R}}(A)}\, z_3^{\rho_{\mathrm{B}}(A)-1}\, z_4^{n-2-\rho_{\mathrm{L}}(A)}$ (at $x=y=1$), which can be shown straightforwardly to be related to generating functions used here by

$$\begin{aligned}
&z_1\, z_2\, z_3\, z_4\, Z'_n(x,y;z_1,z_2,z_3,z_4) = \\
&\quad \widetilde{Z}^{\mathrm{quad}}_n(x,y;z_1,z_2,z_3,z_4) - (z_2z_4)^{n-1}\, Z^{\mathrm{adj}}_{n-1}(x,y;\tfrac{1}{z_4},z_1) - (z_2z_4)^{n-1}\, Z^{\mathrm{adj}}_{n-1}(x,y;\tfrac{1}{z_2},z_3) - \\
&\qquad z_1{z_2}^{n-2}(xz_3)^{n-1}\, \widetilde{Z}^{\mathrm{adj}}_{n-1}(x,y;z_1,\tfrac{1}{z_2}) - z_3{z_4}^{n-2}(xz_1)^{n-1}\, \widetilde{Z}^{\mathrm{adj}}_{n-1}(x,y;z_3,\tfrac{1}{z_4}) + \\
&\qquad\qquad \big((z_2z_4)^{n-1} + x^{2n-3}(z_1z_3)^{n-1}\big)\, Z_{n-2}(x,y). \qquad (51)
\end{aligned}$$

In the matrices given there, functions $\gamma_n(z)$ and $\delta_n(z)$ are used which appear to be related (although this has not been fully confirmed) to the functions used in the matrices here by

$$\begin{aligned}
\gamma_n(z) &= \tfrac{4(n-1)!\,(3n-5)!}{(2n-3)!\,(2n-4)!}\, z^2\, Z_{n-2}(1,1;z) - \tfrac{4(2n-3)!\,(2n-5)!}{(n-3)!\,(3n-5)!}\, (z-1)^2\, Z_n(1,1;z),\\
\delta_n(z) &= \tfrac{4(n-1)!\,(n-4)!\,(3n-5)!\,(3n-8)!}{((2n-4)!)^2\,(2n-5)!\,(2n-7)!}\, z^3\, Z_{n-3}(1,1;z) - \tfrac{9(9n^2-30n+20)}{2n-5}\, z(z-1)^2\, Z_{n-1}(1,1;z). \qquad (52)
\end{aligned}$$

3.4. **Arbitrary bulk parameters x and y.** The case in which the bulk parameters x and y are arbitrary is of primary interest in this paper, and several new results will be presented in Section 4. In this section, the previously-known results for this case, which involve the unrefined, singly-refined and opposite-boundary doubly-refined ASM generating functions, are reviewed.

A determinant formula for the opposite-boundary doubly-refined ASM generating function is

$$Z^{\mathrm{opp}}_n(x,y;z_1,z_2) = \det_{0\le i,j\le n-1}\big(K_n(x,y;z_1,z_2)_{ij}\big), \qquad (53)$$

where

$$\begin{aligned}
&K_n(x,y;z_1,z_2)_{ij} = \\
&\qquad -\delta_{i,j+1} + \begin{cases}
\sum_{k=0}^{\min(i,j+1)} \binom{i-1}{i-k}\binom{j+1}{k} x^k y^{i-k}, & j\le n-3\\
\sum_{k=0}^{i}\sum_{l=0}^{k} \binom{i-1}{i-k}\binom{n-l-2}{k-l} x^k y^{i-k} z_2^{l+1}, & j=n-2\\
\sum_{k=0}^{i}\sum_{l=0}^{k}\sum_{m=0}^{l} \binom{i-1}{i-k}\binom{n-l-2}{k-l} x^k y^{i-k} z_1^m z_2^{l-m}, & j=n-1.
\end{cases} \qquad (54)
\end{aligned}$$

Setting $z_2=1$ or $z_1=z_2=1$ in (53)–(54), and using standard binomial coefficient identities, it follows that determinant formulae for the singly-refined and unrefined ASM generating functions are

$$Z_n(x,y;z) = \det_{0\le i,j\le n-1}\bigl(K_n(x,y;z)_{ij}\bigr), \tag{55}$$

$$Z_n(x,y) = \det_{0\le i,j\le n-1}\bigl(K_n(x,y)_{ij}\bigr), \tag{56}$$

where

$$\begin{aligned} K_n(x,y;z)_{ij} &= K_n(x,y;z,1)_{ij} \\ &= -\delta_{i,j+1} + \begin{cases} \sum_{k=0}^{\min(i,j+1)} \binom{i-1}{i-k}\binom{j+1}{k} x^k y^{i-k}, & j \le n-2 \\ \sum_{k=0}^{i}\sum_{l=0}^{k} \binom{i-1}{i-k}\binom{n-l-1}{k-l} x^k y^{i-k} z^l, & j = n-1, \end{cases} \end{aligned} \tag{57}$$

$$K_n(x,y)_{ij} = K_n(x,y;1,1)_{ij} = -\delta_{i,j+1} + \textstyle\sum_{k=0}^{\min(i,j+1)} \binom{i-1}{i-k}\binom{j+1}{k} x^k y^{i-k}. \tag{58}$$

There are further simplifications if $x = y = 1$. For example,

$$K_n(1,1)_{ij} = -\delta_{i,j+1} + \tbinom{i+j}{i}. \tag{59}$$

The opposite-boundary doubly-refined ASM generating function satisfies

$$\begin{aligned} &(z_1-z_2)\,(z_3-z_4)\,Z_n^{\mathrm{opp}}(x,y;z_1,z_2)\,Z_n^{\mathrm{opp}}(x,y;z_3,z_4) - \\ &\qquad (z_1-z_3)\,(z_2-z_4)\,Z_n^{\mathrm{opp}}(x,y;z_1,z_3)\,Z_n^{\mathrm{opp}}(x,y;z_2,z_4) + \\ &\qquad\qquad (z_1-z_4)\,(z_2-z_3)\,Z_n^{\mathrm{opp}}(x,y;z_1,z_4)\,Z_n^{\mathrm{opp}}(x,y;z_2,z_3) = 0, \end{aligned} \tag{60}$$

and it can be expressed in terms of singly-refined and unrefined ASM generating functions as

$$\begin{aligned} (z_1-z_2)\,Z_n^{\mathrm{opp}}(x,y;z_1,z_2)\,Z_{n-1}(x,y) &= (z_1-1)\,z_2\,Z_n(x,y;z_1)\,Z_{n-1}(x,y;z_2) - \\ &\qquad z_1\,(z_2-1)\,Z_{n-1}(x,y;z_1)\,Z_n(x,y;z_2). \end{aligned} \tag{61}$$

The two previous identities are essentially equivalent, since (61) can be obtained from (60) by setting $z_3 = 1$ and $z_4 = 0$, and then applying relations from (11) and (13), while (60) can be obtained from (61) by expressing each of the six cases of $(z_i - z_j)Z_n^{\mathrm{opp}}(x,y;z_i,z_j)$ in terms of singly-refined and unrefined ASM generating functions, and then checking that the resulting expression for the LHS of (60) vanishes.

The formula (53) was derived by Behrend, Di Francesco and Zinn-Justin [16, Eqs. (21)–(22)], as part of the proof of an equality [16, Thm. 1] between the opposite-boundary doubly-refined ASM generating function and a four-parameter generating function for descending plane partitions. (See also Section 3.12.) The formula (55) was derived by Behrend, Di Francesco and Zinn-Justin [17, Eqs. (97)–(98)], prior to the derivation of (53), as part of the proof of an equality [17, Thm. 1] (conjectured previously by Mills, Robbins and Rumsey [103, Conj. 3]) between the singly-refined ASM generating function and a three-parameter generating function for descending plane partitions.

For an alternative version of (53), involving a transformation of the matrix $K_n(x,y;z_1,z_2)$, see Behrend, Di Francesco and Zinn-Justin [16, Eqs. (65)–(66)], and for alternative versions of (55) or (56), involving transformations of the matrices $K_n(x,y;z)$ or $K_n(x,y)$, see

Behrend, Di Francesco and Zinn-Justin [17, Props. 1 & 4, Eqs. (65), (66), (87) & (88), & Sec. 4.1]. For additional information on determinants closely related to those of (55)–(56), including formulae for their evaluation or factorization in certain special cases, see, for example, Andrews [2], Andrews and Stanton [5], Ciucu, Eisenkölbl, Krattenthaler and Zare [38, Thms. 10–13], Ciucu and Krattenthaler [39], Colomo and Pronko [41, Eqs. (23)–(24)], [42, Eqs. (4.3)–(4.7)], de Gier [53, Sec. 2.1], Gessel and Xin [76, Sec. 5], Krattenthaler [88, e.g., Thms. 25–37], [89, Sec. 5.5], Lalonde [94, Thm. 3.1], Mills, Robbins and Rumsey [102], [103, p. 346], [105, Secs. 3–4], Robbins [124, Sec. 2], and Rosengren [128].

The case of (61) with $x = y = 1$ (as given in (43)) was first obtained by Stroganov [140, Sec. 5], and this relation for arbitrary x and y was first obtained, as an identity involving one- and two-point boundary correlation functions for the six-vertex model with DWBC, by Colomo and Pronko [44, Eq. (5.32)], [46, Eq. (3.32)]. An alternative proof of (61) was given by Behrend, Di Francesco and Zinn-Justin [16, Sec. 5]. This identity will also be obtained from Theorem 1 of this paper, as Corollary 4, and it will be found to be a special case of the identity (71) given in Section 3.5.

3.5. **ASM enumeration involving statistics associated with several rows or several columns.**

In this section, a generating function involving statistics associated with several rows (or several columns) of ASMs is defined, and (in (71)) a result is stated which, for the case of particular assignments of certain parameters, provides a determinantal expression for this generating function in terms of singly-refined and unrefined ASM generating functions. A derivation of this result will be given in Section 5.9.

For $1 \le i, j \le n$ and $A \in \mathrm{ASM}(n)$, define ASM statistics, associated with row i or column j of A, as

$$\begin{aligned}
\nu^{\mathrm{row}\,i}(A) &= \bigl|\{1 \le j \le n \mid \textstyle\sum_{i'=1}^{i-1} A_{i'j} = \sum_{j'=1}^{j} A_{ij'}\}\bigr|,\\
\nu^{\mathrm{col}\,j}(A) &= \bigl|\{1 \le i \le n \mid \textstyle\sum_{i'=1}^{i-1} A_{i'j} = \sum_{j'=1}^{j} A_{ij'}\}\bigr|,\\
\mu^{\mathrm{row}\,i}(A) &= \text{number of } -1\text{'s in row } i \text{ of } A,\\
\mu^{\mathrm{col}\,j}(A) &= \text{number of } -1\text{'s in column } j \text{ of } A.
\end{aligned} \tag{62}$$

It follows from the defining properties of ASMs that if, for an ASM A, $\sum_{i'=1}^{i-1} A_{i'j} = \sum_{j'=1}^{j} A_{ij'}$ (or equivalently $\sum_{i'=1}^{i} A_{i'j} = \sum_{j'=1}^{j-1} A_{ij'}$), then $A_{ij} = 0$. Hence, $\nu^{\mathrm{row}\,i}(A)$ and $\nu^{\mathrm{col}\,j}(A)$ can be regarded as the numbers of certain 0's in row i and column j, respectively, of A. As will be seen in Section 5.2, for the configuration of the six-vertex model with DWBC which corresponds to A, $\nu^{\mathrm{row}\,i}(A)$ and $\nu^{\mathrm{col}\,j}(A)$ are simply the numbers of vertex configurations of types (1) and (2) in row i and column j, respectively, of the grid.

It can also be seen that the statistics $\nu^{\mathrm{row}\,i}(A)$ and $\nu^{\mathrm{col}\,j}(A)$ can be written in various other ways, for example as

$$\begin{aligned}
\nu^{\mathrm{row}\,i}(A) &= \textstyle\sum_{j=1}^{n}\bigl(\sum_{i'=1}^{i-1}\sum_{j'=1}^{j} A_{i'j}A_{ij'} + \sum_{i'=i}^{n}\sum_{j'=j+1}^{n} A_{i'j}A_{ij'}\bigr),\\
\nu^{\mathrm{col}\,j}(A) &= \textstyle\sum_{i=1}^{n}\bigl(\sum_{i'=1}^{i-1}\sum_{j'=1}^{j} A_{i'j}A_{ij'} + \sum_{i'=i}^{n}\sum_{j'=j+1}^{n} A_{i'j}A_{ij'}\bigr).
\end{aligned} \tag{63}$$

It can be checked easily that, for any $A \in \mathrm{ASM}(n)$, the statistics of (3)–(4) and those of (62) are related by

$$\begin{gathered}
\textstyle\sum_{i=1}^{n} \nu^{\mathrm{row}\,i}(A) = \sum_{j=1}^{n} \nu^{\mathrm{col}\,j}(A) = 2\nu(A),\\
\nu^{\mathrm{row}\,1}(A) = \rho_{\mathrm{T}}(A), \quad \nu^{\mathrm{col}\,n}(A) = \rho_{\mathrm{R}}(A), \quad \nu^{\mathrm{row}\,n}(A) = \rho_{\mathrm{B}}(A), \quad \nu^{\mathrm{col}\,1}(A) = \rho_{\mathrm{L}}(A),\\
\textstyle\sum_{i=2}^{n-1} \mu^{\mathrm{row}\,i}(A) = \sum_{j=2}^{n-1} \mu^{\mathrm{col}\,j}(A) = \mu(A),\\
\mu^{\mathrm{row}\,1}(A) = \mu^{\mathrm{col}\,n}(A) = \mu^{\mathrm{row}\,n}(A) = \mu^{\mathrm{col}\,1}(A) = 0,
\end{gathered} \tag{64}$$

and that, for any $A \in \mathrm{ASM}(n)$ and $1 \le i \le n$, the statistics of (62) behave under the operations of transposition or anticlockwise quarter-turn rotation of ASMs according to

$$\begin{aligned}
&\nu^{\mathrm{row}\,i}(A) = \nu^{\mathrm{col}\,i}(A^T) = n-1-\nu^{\mathrm{col}\,i}(A^Q) - 2\mu^{\mathrm{row}\,i}(A),\\
&\mu^{\mathrm{row}\,i}(A) = \mu^{\mathrm{col}\,i}(A^T) = \mu^{\mathrm{col}\,i}(A^Q),
\end{aligned} \tag{65}$$

using the same notation as in (6).

For any $0 \le m \le n$ and $1 \le k_1 < \ldots < k_m \le n$, define an ASM generating function associated with rows (or, as will be seen in (68), columns) $k_1, \ldots, k_m$ as

$$Z_n^{k_1,\ldots,k_m}(x,y;z_1,\ldots,z_m;w_1,\ldots,w_m) = \\ \sum_{A\in\mathrm{ASM}(n)} x^{\nu(A)}\, y^{\mu(A)-\sum_{i=1}^m \mu^{\mathrm{row}\,k_i}(A)} \prod_{i=1}^m z_i^{\nu^{\mathrm{row}\,k_i}(A)}\, w_i^{\mu^{\mathrm{row}\,k_i}(A)}, \tag{66}$$

for indeterminates x, y, $z_1, \ldots, z_m$ and $w_1, \ldots, w_m$.

It can be seen that

$$Z_n^{k_1,\ldots,k_m}(x,y;z_1,\ldots,z_{i-1},1,z_{i+1},\ldots,z_m;w_1,\ldots,w_{i-1},y,w_{i+1},\ldots,w_m) = \\ Z_n^{k_1,\ldots,k_{i-1},k_{i+1},\ldots,k_m}(x,y;z_1,\ldots,z_{i-1},z_{i+1},\ldots,z_m;w_1,\ldots,w_{i-1},w_{i+1},\ldots,w_m), \tag{67}$$

and that $Z_n^{k_1,\ldots,k_m}(x,y;z_1,\ldots,z_m;w_1,\ldots,w_m)$ is the unrefined ASM generating function $Z_n(x,y)$ for $m=0$, the singly-refined ASM generating function $Z_n(x,y;z_1)$ for $m=1$ and $k_1=1$ or $k_1=n$, and the opposite-boundary doubly-refined ASM generating function $Z_n^{\mathrm{opp}}(x,y;z_1,z_2)$ for $m=2$, $k_1=1$ and $k_2=n$.

Using (65), it also follows that

$$Z_n^{k_1,\ldots,k_m}(x,y;z_1,\ldots,z_m;w_1,\ldots,w_m) = \\ \sum_{A\in\mathrm{ASM}(n)} x^{\nu(A)}\, y^{\mu(A)-\sum_{j=1}^m \mu^{\mathrm{col}\,k_j}(A)} \prod_{j=1}^m z_j^{\nu^{\mathrm{col}\,k_j}(A)}\, w_j^{\mu^{\mathrm{col}\,k_j}(A)} = \\ x^{n(n-1)/2}\,(z_1\ldots z_m)^{n-1}\, Z_n^{k_1,\ldots,k_m}\bigl(\tfrac{1}{x},\tfrac{y}{x};\,\tfrac{1}{z_1},\ldots,\tfrac{1}{z_m};\,\tfrac{w_1}{xz_1^2},\ldots,\tfrac{w_m}{xz_m^2}\bigr). \tag{68}$$

Now define, for $0 \le m \le n$, a further function

$$X_n(x,y;z_1,\ldots,z_m) = \begin{cases} Z_n(x,y), & m=0 \\ \dfrac{\det_{1\le i,j\le m}\bigl(z_i^{\,j-1}\,(z_i-1)^{m-j}\,Z_{n-j+1}(x,y;z_i)\bigr)}{\prod_{1\le i<j\le m}(z_i-z_j)\,\prod_{i=1}^{m-1} Z_{n-i}(x,y)}, & 1\le m\le n. \end{cases} \tag{69}$$

This definition is based on a definition of Colomo and Pronko [47, Eq. 6.6], [49, Eq. (4.18)] of a closely-related function for the six-vertex model with DWBC.

Some properties of $X_n(x,y;z_1,\ldots,z_m)$, which follow straightforwardly from (69) and the latter equations of (11), (12) and (13), are

$$\begin{gathered} X_n(x,y;z_1,\ldots,z_m) \text{ is symmetric in } z_1,\ldots,z_m, \\ X_n(x,y;z_1,\ldots,z_m) = x^{n(n-1)/2}\,(z_1\ldots z_m)^{n-1}\,X_n\bigl(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_1},\ldots,\tfrac{1}{z_m}\bigr), \\ X_n(x,y;z_1,\ldots,z_{i-1},1,z_{i+1},\ldots,z_m) = X_n(x,y;z_1,\ldots,z_{i-1},z_{i+1},\ldots,z_m), \\ X_n(x,y;z_1,\ldots,z_{i-1},0,z_{i+1},\ldots,z_m) = X_{n-1}(x,y;z_1,\ldots,z_{i-1},z_{i+1},\ldots,z_m). \end{gathered} \tag{70}$$

The main result of this section is that, for any $0 \le m \le n$ and $1 \le k_1 < \ldots < k_m \le n$,

$$Z_n^{k_1,\ldots,k_m}(x,y;z_1,\ldots,z_m;xz_1^2+(y-x-1)z_1+1,\ldots,xz_m^2+(y-x-1)z_m+1) = X_n(x,y;z_1,\ldots,z_m). \tag{71}$$

A derivation of (71), based on a result of Colomo and Pronko [47, Eq. (6.8)], [48, Eq. (A.13)] will be given in Section 5.9.

Note that, using (67) and the third property of (70), the cases of (71) with $m<n$ all follow from the case $m=n$.

It can now be seen, using (66) and (71), that the function (69) is a polynomial in x, y and $z_1,\ldots,z_m$ with integer coefficients.

It follows from (71) that, for fixed n and m, the LHS of (71) is independent of $k_1,\ldots,k_m$. This implies, for example, that the singly-refined and opposite-boundary doubly-refined ASM generating functions can be written, for any $1 \le k \le n$ and $1 \le k_1 < k_2 \le n$, as

$$\begin{aligned} Z_n(x,y;z) &= Z_n^k(x,y;z;xz^2+(y-x-1)z+1), \\ Z_n^{\mathrm{opp}}(x,y;z_1,z_2) &= Z_n^{k_1,k_2}(x,y;z_1,z_2;xz_1^2+(y-x-1)z_1+1,xz_2^2+(y-x-1)z_2+1). \end{aligned} \tag{72}$$

It can also be seen that (71) contains the identity (61), satisfied by the opposite-boundary doubly-refined ASM generating function, as the case $m=2$, $k_1=1$ and $k_2=n$.

It will be shown in Section 5.10 that the function (69) at $x=y=1$ can be expressed as

$$X_n(1,1;z_1,\ldots,z_m) = 3^{-n(n-1)/2}\,(-q)^{m(n-1)}\,\bigl((z_1+q)\ldots(z_m+q)\bigr)^{n-1}\times \\ s_{(n-1,n-1,\ldots,2,2,1,1)}\Bigl(\tfrac{qz_1+1}{z_1+q},\ldots,\tfrac{qz_m+1}{z_m+q},\underbrace{1,\ldots,1}_{2n-m}\Bigr)\Big|_{q=e^{\pm 2\pi i/3}}, \tag{73}$$

where this uses the same notation as, and can be seen to generalize, the identity (45).

3.6. **ASMs with several rows or columns closest to two opposite boundaries prescribed.** In this section, the enumeration of ASMs with prescribed configurations of several rows or columns closest to two opposite boundaries (but not involving any further statistics) is discussed.

Let K_1 and K_2 be subsets of $\{1,\ldots,n\}$ with $|K_1|+|K_2|\le n$, and let A_{K_1} and A_{K_2} be matrices with n columns and $|K_1|$ or $|K_2|$ rows respectively, in which each entry is 0, 1 or -1, along each row and column the nonzero entries alternate in sign, in each row the entries sum to 1, in column j of A_{K_1} the entries sum to 1 if $j\in K_1$ and the entries sum to 0 with the first nonzero entry (if there is one) being 1 if $j\notin K_1$, and in column j of A_{K_2} the entries sum to 1 if $j\in K_2$ and the entries sum to 0 with the first nonzero entry (if there is one) being -1 if $j\notin K_2$. (For example, the entries of these matrices could be taken as simply $(A_{K_1})_{ij}=\delta_{j,K_{1,i}}$ and $(A_{K_2})_{ij}=\delta_{j,K_{2,i}}$, where $K_{1,i}$ and $K_{2,i}$ are the ith smallest elements of K_1 and K_2, respectively.) For a fixed choice of matrices A_{K_1} and A_{K_2} which satisfy these conditions, define

$$\begin{aligned}\mathcal{A}_{n,K_1,K_2} = \text{number of } n\times n \text{ ASMs whose first } |K_1| \text{ rows are given by } A_{K_1},\\ \text{and last } |K_2| \text{ rows are given by } A_{K_2},\end{aligned}\tag{74}$$

where it can be checked easily that $\mathcal{A}_{n,K_1,K_2}$ is independent of the choice of A_{K_1} and A_{K_2}. (Note that if $K_1=\emptyset$ or $K_2=\emptyset$, then there is no restriction on the first or last rows, respectively, of the ASMs in (74).) Thus, $\mathcal{A}_{n,K_1,K_2}$ is the number of $n\times n$ ASMs in which certain rows or columns closest to two opposite boundaries are prescribed.

Alternatively, $\mathcal{A}_{n,K_1,K_2}$ can be written, without reference to A_{K_1} and A_{K_2}, as the number of $(n-|K_1|-|K_2|)\times n$ matrices in which each entry is 0, 1 or -1, along each row and column the nonzero entries alternate in sign, in each row the entries sum to 1, and in column j the entries sum to 1 if $j\in\{1,\ldots,n\}\setminus(K_1\cup K_2)$, the entries sum to -1 if $j\in K_1\cap K_2$, the entries sum to 0 with the first nonzero entry (if there is one) being -1 if $j\in K_1\setminus K_2$, and the entries sum to 0 with the first nonzero entry (if there is one) being 1 if $j\in K_2\setminus K_1$.

It can be seen that

$$\mathcal{A}_{n,K_1,K_2}=\mathcal{A}_{n,K_2,K_1}=\mathcal{A}_{n,\{n+1-k|k\in K_1\},\{n+1-k|k\in K_2\}}\tag{75}$$

and that

$$\mathcal{A}_{n,\emptyset,\emptyset}=\mathcal{A}_n,\quad \mathcal{A}_{n,\{k+1\},\emptyset}=\mathcal{A}_{n,k},\quad \mathcal{A}_{n,\{k_1+1\},\{n-k_2\}}=\mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2},\tag{76}$$

for any $0\le k,k_1,k_2\le n-1$.

It can also be seen that $\mathcal{A}_{n,\{1,\ldots,n\}\setminus K_1,\emptyset}$ or $\mathcal{A}_{n,\{1,\ldots,n\}\setminus K_2,\emptyset}$ give the numbers of possible matrices A_{K_1} or A_{K_2}, respectively, which satisfy the conditions outlined above. Furthermore, for a fixed set K of positive integers, $\mathcal{A}_{n,\{1,\ldots,n\}\setminus K,\emptyset}$ is independent of n, for $n\ge\max(K)$, with an operator formula for this number having been obtained by Fischer [67, Thm. 1]. Some related functions for the six-vertex model with DWBC have been studied by Colomo and Pronko [49].

The numbers $\mathcal{A}_{n,K_1,K_2}$ have been studied by Fischer [70, 71], Fischer and Romik [72], and Karklinsky and Romik [83]. Results which have been obtained include linear relations between the numbers $\mathcal{A}_{n,K,\emptyset}$ with $|K| = 2$ and the opposite-boundary doubly-refined ASM numbers (Karklinsky and Romik [83, Eq. (7)], and Fischer [70, Eq. (5.3)]), explicit formulae for $\mathcal{A}_{n,K,\emptyset}$ with $|K| = 2$ (Karklinsky and Romik [83, Thm. 1], and Fischer [70, Eq. (1.3)]), and an expression for $\mathcal{A}_{n,K,\{k\}}$ with $|K| = 2$ in terms of numbers of $n \times n$ ASMs with prescribed configurations on three boundaries (Fischer [71, Thm. 1]). Also, related numbers have been defined (Fischer [70, Eq. (1.2)], [71, Eq. (2.9)]), and have been shown to give the numbers $\mathcal{A}_{n,K_1,K_2}$ in certain cases (Fischer [70, Thm. 1]), and to satisfy certain linear relations (Fischer [70, Eq. (5.1) & p. 253, first equation], [71, Eqs. (1.5) & (1.6)]).

3.7. **The cases $Z_n(1,3)$ and $Z_n(1,3;z)$.**

Results involving $Z_n(1,3)$ and $Z_n(1,3;z)$, some of which were previously conjectured by Mills, Robbins and Rumsey [103, Conj. 6–7], [104, Conj. 5], and which correspond to the so-called 3-enumeration of ASMs, have been obtained by Colomo and Pronko [43], [45, Secs. 4.4 & 5.4], [46, Sec. 4.3], Kuperberg [92, Thm. 2], [93, Thm. 3], Okada [111, Thm. 2.4(1), fourth eq.], and Stroganov [138].

3.8. **The case $Z_n(1,1;-1)$.**

It was observed by Di Francesco [57, Eqs. (2.7)–(2.8)], and shown explicitly by Williams [144, Thm. 4], that

$$Z_n(1,1;-1) = \mathcal{V}_n^2, \tag{77}$$

where $\mathcal{V}_n$ is the number of vertically-symmetric $n \times n$ ASMs (i.e., ASMs A with $A_{ij} = A_{i,n+1-j}$). Note that for n even, it can be seen easily (for example, using the second equation of (32)) that both sides of the equation are zero. For n odd, $\mathcal{V}_n = \prod_{i=1}^{(n-1)/2}(6i-2)!/(n+2i-1)!$, as conjectured by Robbins [123, 124], and first shown by Kuperberg [93, Thm. 2, second equation]. It is known that $\mathcal{V}_n$ is also the number of descending plane partitions invariant under a certain operation (see Behrend, Di Francesco and Zinn-Justin [17, Sec. 4.2 & Eq. (102)]), and the number of totally symmetric self-complementary plane partitions invariant under a certain operation (see Ishikawa [80, Thm. 7.11(i)]), and it has been conjectured by Di Francesco [57, Eq. (2.8)], [58, Eq. (4.5)] that $\mathcal{V}_n^2$ (up to sign) and $\mathcal{V}_n^4$ are obtained when a parameter is set to -1 in certain generating functions for totally symmetric self-complementary and cyclically symmetric transpose-complementary plane partitions.

3.9. **ASMs with a fixed number of generalized inversions.**

Expressions for the coefficients of x^p in $Z_n(x,1)$, i.e., for the number of $n \times n$ ASMs with p generalized inversions, can be obtained using a result of Behrend [15, Cor. 14], and are given by Behrend, Di Francesco and Zinn-Justin [17, Eqs. (24)–(25)]. Some particular cases of p, for ASMs with fixed values of certain further statistics, are also considered by Behrend, Di Francesco and Zinn-Justin [16, Sec. 7.1], [17, pp. 337–338].

3.10. **ASMs with a fixed number of -1's.**

For the case of $Z_n(1,y)$, an expression for the coefficient of y^m, i.e., for the number of $n \times n$ ASMs with m -1s, has been obtained by Cori, Duchon and Le Gac [50, 99] (and, for $m = 1,2$, by Aval [6, Prop. 4]), a certain factorization property (previously conjectured by Mills, Robbins and Rumsey [103, Conj. 4 & Conj. 5], [105, pp. 50 & 54] and Robbins [124, Sec. 2]) has been established by Kuperberg [92, Thm. 3], [93, Thm. 4, first two eqs.], and a certain operator formula containing a parameter which corresponds to the number of -1's in an ASM has been obtained by Fischer [69, Thm. 1].

Results involving the coefficient of y^1 in $Z_n(x,y;z)$, i.e., for a certain generating function for $n\times n$ ASMs with a single -1, have been obtained by Lalonde [94, 96].

3.11. **Objects in simple bijection with ASMs.** Simple bijections are known between ASMs and various other combinatorial objects. Some examples, all of which are reviewed by Propp [113, Secs. 2–4 & 7], are bijections between $\mathrm{ASM}(n)$ and sets of $(n+1)\times(n+1)$ corner-sum matrices, $(n+1)\times(n+1)$ height-function matrices, 3-colorings of an $(n+1)\times(n+1)$ grid with certain boundary conditions, monotone (or Gog) triangles with bottom row $1,\ldots,n$, configurations of the six-vertex model on an $n\times n$ grid with DWBC, and fully packed loop configurations on an $n\times n$ grid with certain boundary conditions. Some further examples are a bijection between $\mathrm{ASM}(n)$ and the set of sets of n osculating paths on an $n\times n$ grid in which all paths start and end on two adjacent boundaries (see, for example, Behrend [15, Secs. 2–4], Bousquet-Mélou and Habseiger [21, pp. 68–69], Bressoud [22, pp. 226–227], or Eğecioğlu, Redmond and Ryavec [62, pp. 35–36]), and a bijection between $\mathrm{ASM}(n)$ and the set of alternating paths for any fixed fully packed loop configuration on an $n\times n$ grid (see Ng [110, Prop. 3.1]).

In each of these cases, the statistics for the other combinatorial object which correspond, under the bijection, to the ASM statistics (3)–(4) or (62) can be obtained straightforwardly. For example, this will be done for configurations of the six-vertex model with DWBC in (111). Accordingly, all of the results reviewed or obtained in this paper for $\mathrm{ASM}(n)$ with the statistics (3)–(4) or (62) can alternatively be expressed in terms of any of these other combinatorial objects and their associated statistics.

In the case of the six-vertex model with DWBC, ASM generating functions are closely related to certain partition functions, expectation values, probabilities or correlation functions for the model, and the results of this paper could alternatively be expressed in terms of the latter quantities. For example, the quadruply-refined ASM generating function is closely associated with a certain four-point boundary correlation function for the six-vertex model with DWBC. For studies of various correlation functions, and related quantities, for the six-vertex model with DWBC, see, for example, Bogoliubov, Kitaev and Zvonarev [19], Bogoliubov, Pronko and Zvonarev [20], Colomo and Pronko [44, 46, 47, 49], Foda and Preston [73], and Motegi [106, 107].

3.12. **Descending plane partitions.** Descending plane partitions are certain combinatorial objects first defined by Andrews [2, 3]. It was shown by Behrend, Di Francesco and Zinn-Justin [16, Thm. 1] that the opposite-boundary doubly-refined ASM generating function $Z_n^{\mathrm{opp}}(x,y;z_1,z_2)$ is equal to the generating function for descending plane partitions with largest part at most n, in which the statistics associated with x, y, z_1 and z_2 are, respectively, the number of nonspecial parts, the number of special parts (as first defined by Mills, Robbins and Rumsey [103, p. 344]), the number of n's, and the number of $(n-1)$'s plus the number of rows of length $n-1$, in a descending plane partition. (Note that no further statistics for descending plane partitions are currently known which, together with the previous four statistics, lead to generating functions which are equal to the adjacent-boundary doubly-refined, triply-refined or quadruply-refined ASM generating functions, and that no general bijection is currently known between $\mathrm{ASM}(n)$ and the set of descending plane partitions with largest part at most n.)

It follows that certain known results for descending plane partitions correspond to results involving the opposite-boundary doubly-refined, singly-refined or unrefined ASM generating functions. For example, certain such results were outlined in Section 3.4.

Results are also known for the enumeration of descending plane partitions with a prescribed sum of parts. In particular, it follows from work of Mills, Robbins and Rumsey [102, Sec. 5] that, for $0 \le k \le n-1$,

$$\begin{aligned}\textstyle\sum_{D\in\mathrm{DPP}(n)} q^{|D|} &= \textstyle\prod_{i=0}^{n-1} \frac{[3i+1]_q!}{[n+i]_q!},\\ \textstyle\sum_{D\in\mathrm{DPP}(n,k)} q^{|D|} &= \textstyle\frac{q^{kn}\,[n+k-1]_q!\,[2n-k-2]_q!}{[k]_q!\,[n-k-1]_q!\,[2n-2]_q!}\,\prod_{i=0}^{n-2} \frac{[3i+1]_q!}{[n+i-1]_q!},\end{aligned}\tag{78}$$

where $\mathrm{DPP}(n)$ denotes the set of descending plane partitions with largest part at most n, $\mathrm{DPP}(n,k)$ denotes the set of elements of $\mathrm{DPP}(n)$ with exactly k parts equal to n, $|D|$ denotes the sum of parts of a descending plane partition D, and $[n]_q!$ is the usual q-factorial of n. As expected, each RHS of (78) reduces to each RHS of (33)–(34) for $q=1$. However, no ASM statistic is currently known which corresponds to the sum of parts of a descending plane partition, and leads to generating functions matching those of (78).

For further information, and related results or conjectures, regarding descending plane partitions, see, for example, Andrews [2, 3], Ayyer [10], Behrend, Di Francesco and Zinn-Justin [16, 17], Bresssoud [22], Bressoud and Propp [25], Krattenthaler [90], Lalonde [94, 95, 96], Mills, Robbins and Rumsey [102, 103, 105], Robbins [123, 124], and Striker [135].

3.13. **Totally symmetric self-complementary plane partitions.** Totally symmetric self-complementary plane partitions are certain combinatorial objects first defined by Mills, Robbins and Rumsey [104, Sec. 1], and Stanley [131, Sec. 2]. It was shown by Fonseca and Zinn-Justin [75], following conjectures of Mills, Robbins and Rumsey [104, Conj. 2 & 3], that the opposite-boundary doubly-refined ASM generating function with both bulk parameters set to 1, $Z_n^{\mathrm{opp}}(1,1;z_1,z_2)$, is equal to a generating function for totally symmetric self-complementary plane partitions in a $2n\times 2n\times 2n$ box in which the statistics associated with z_1 and z_2 can be certain pairs from among several statistics defined by Doran [61, Sec. 7], and Mills, Robbins and Rumsey [104, Sec. 3]. The singly-refined case of this result was obtained earlier by Razumov, Stroganov and Zinn-Justin [122, Sec. 5.5]. (Note that no statistics for totally symmetric self-complementary plane partitions are currently known which have the same enumerative behavior as the number of generalized inversions or the number of -1's in an ASM, and that no general bijection is currently known between $\mathrm{ASM}(n)$ and the set of totally symmetric self-complementary plane partitions in a $2n\times 2n\times 2n$ box.)

It follows that certain known results for totally symmetric self-complementary plane partitions correspond to results involving the ASM generating functions $Z_n^{\mathrm{opp}}(1,1;z_1,z_2)$ or $Z_n(1,1;z)$, or the ASM numbers $\mathcal{A}^{\mathrm{opp}}_{n,k_1,k_2}$, $\mathcal{A}_{n,k}$ or $\mathcal{A}_n$. For example, for $Z_n^{\mathrm{opp}}(1,1;z_1,z_2)$, $Z_n(1,1;z)$ or $\mathcal{A}_n$, Pfaffian expressions follow from results of Ishikawa [79, Thms. 1.2 & 1.4, & Sec. 7] and Stembridge [133, Thm. 8.3], constant-term expressions follow from results of Ishikawa [79, Sec. 8], Krattenthaler [87, Thm.] and Zeilberger [145, Sublems. 1.1 & 1.2], and integral expressions follow from results of Fonseca and Zinn-Justin [75, Eqs. (4.9) & (4.14)]. Several of these results include a certain ASM statistic, first introduced by Mills, Robbins and Rumsey [104, p. 287], but not considered in this paper.

For the case of adjacent-boundary doubly-refined ASM enumeration, it has been conjectured by Cheballah [34, Conj. 4.3.1] that $\mathcal{A}^{\mathrm{adj}}_{n,k_1,k_2}$ equals the number of totally symmetric self-complementary plane partitions in a $2n\times2n\times2n$ box for which a certain pair of statistics have values k_1 and k_2.

For further information, and related results or conjectures, regarding totally symmetric self-complementary plane partitions, see, for example, Andrews [4], Ayyer, Cori and Gouyou-Beauchamps [11], Bressoud [22, Sec. 6.2], Bressoud and Propp [25], Cheballah [34], Cheballah and Biane [35], Di Francesco [57, 58], Doran [61], Fonseca [74, Secs. 3–4], Fonseca and Zinn-Justin [75], Ishikawa [79, 80], Krattenthaler [87], Mills, Robbins and Rumsey [104], Robbins [123], Stanley [131], Stembridge [133, Sec. 8], Striker [134, 136], Zeilberger [145], Zinn-Justin [149], and Zinn-Justin and Di Francesco [150].

3.14. **Loop models.** Numbers of certain ASMs, or plane partitions, have been found to appear also as appropriately normalized entries (or sums of such entries) of particular eigenvectors associated with certain cases of integrable loop models (or associated quantum spin chains). A wide variety of such cases are known, and these often involve ASMs with prescribed link patterns of associated fully packed loop configurations, with prescribed values of some of the statistics of (3)-(4) or of related statistics or parameters, or subject to invariance under certain symmetry operations.

Of the many papers which study the confirmed or conjectured appearances of numbers of ASMs in such contexts, a few examples are Batchelor, de Gier and Nienhuis [13], Cantini and Sportiello [31, 32], Di Francesco [55, 56], Di Francesco and Zinn-Justin [59], Pasquier [112], Razumov and Stroganov [115, 116, 118], and Stroganov [137]. For reviews of some of these matters, see, for example, de Gier [51, 52, 53], or Zinn-Justin [149].

3.15. **ASMs invariant under symmetry operations.** This paper is focussed on the enumeration of ASMs with prescribed values of the bulk statistics (3) and boundary statistics (4), but with no other conditions applied. However, other studies have focussed on the enumeration of ASMs which are invariant under certain symmetry operations, or subject to related conditions, and which in some cases also have prescribed values of some of the statistics of (3)-(4). For results of this type (some of which remain conjectural), see, for example, Aval and Duchon [8, 9], Bousquet-Mélou and Habseiger [21], Bressoud [22, Sec. 6.1], Kuperberg [93], Okada [111], Robbins [123, 124], Razumov and Stroganov [117, 119, 120], Stanley [130], and Stroganov [139, 141].

Numbers of certain such ASMs have also been related (again, in some cases, only conjecturally) to numbers of certain descending plane partitions (see, for example, de Gier, Pyatov and Zinn-Justin [54, Prop. 3, first equation], and Mills, Robbins and Rumsey [103, Conj. 3S]), numbers of certain totally-symmetric self-complementary plane partitions (see, for example, Ishikawa [79, 80], and Mills, Robbins and Rumsey [104, Conjs. 4 & 6]), or appropriately normalized entries of eigenvectors associated with certain cases of loop models (see, for example, the references given in Section 3.14).

4. New Results

In this section, the main result of this paper is stated, and some of its implications are derived and discussed. The proof of the result will be deferred to Section 5.6.

4.1. Main result.

The primary result of this paper is as follows.

Theorem 1. *The quadruply-refined ASM generating function satisfies*

$$
\begin{aligned}
&y(z_4-z_2)(z_1-z_3)\,Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)\,Z_{n-2}(x,y) = \\
&\quad\big((z_1-1)(z_2-1)-yz_1z_2\big)\big((z_3-1)(z_4-1)-yz_3z_4\big)\,Z_{n-1}^{\mathrm{adj}}(x,y;z_4,z_1)\,Z_{n-1}^{\mathrm{adj}}(x,y;z_2,z_3)\,- \\
&\quad\big(x(z_4-1)(z_1-1)-y\big)\big(x(z_2-1)(z_3-1)-y\big)\,z_1z_2z_3z_4\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,z_2)\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_3,z_4)\,- \\
&\qquad(z_2-1)(z_3-1)\big((z_4-1)(z_1-1)-yz_4z_1\big)\,Z_{n-1}^{\mathrm{adj}}(x,y;z_4,z_1)\,Z_{n-2}(x,y)\,+ \\
&\quad(z_3-1)(z_4-1)\big(x(z_1-1)(z_2-1)-y\big)\,z_1z_2\,(xz_3z_4)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,z_2)\,Z_{n-2}(x,y)\,- \\
&\qquad(z_4-1)(z_1-1)\big((z_2-1)(z_3-1)-yz_2z_3\big)\,Z_{n-1}^{\mathrm{adj}}(x,y;z_2,z_3)\,Z_{n-2}(x,y)\,+ \\
&\quad(z_1-1)(z_2-1)\big(x(z_3-1)(z_4-1)-y\big)\,z_3z_4\,(xz_1z_2)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_3,z_4)\,Z_{n-2}(x,y)\,+ \\
&\qquad\qquad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\big(1-(x^2z_1z_2z_3z_4)^{n-1}\big)\,Z_{n-2}(x,y)^2.
\end{aligned}
\tag{79}
$$

The identity (79) holds for each $n \geq 3$. Furthermore, if $Z_0(x,y)$ is taken to be 1, then it can be seen, using $Z_1^{\mathrm{adj}}(x,y;z_1,z_2) = \widetilde{Z}_1^{\mathrm{adj}}(x,y;z_1,z_2) = 1$ and $Z_2^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4) = 1 + xz_1z_2z_3z_4$ (from (8)–(10)), that (79) also holds for $n = 2$.

The proof of (79), which will be given in Section 5.6, will involve using a relation between a certain generalized ASM generating function and the partition function of the six-vertex model with DWBC, and then combining the Desnanot–Jacobi determinant identity with the Izergin–Korepin formula for this partition function.

An alternative form of (79) will be obtained, and some further related results will be discussed, in Section 5.7.

It can be seen that (79) enables $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ to be obtained recursively, using the initial conditions (from (8)) $Z_1^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4) = 1$ and $Z_2^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4) = 1 + xz_1z_2z_3z_4$, and the definitions (from (9)–(10)) $Z_n^{\mathrm{adj}}(x,y;z_1,z_2) = Z_n^{\mathrm{quad}}(x,y;z_1,1,1,z_2)$, $\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2) = Z_n^{\mathrm{quad}}(x,y;z_1,z_2,1,1)$ and $Z_n(x,y) = Z_n^{\mathrm{quad}}(x,y;1,1,1,1)$. Accordingly, in this sense, $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$, and all of the ASM generating functions of (9)–(10) which are defined in terms of $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$, are determined by (79).

Note, however, that if the generating functions are computed recursively in this way, then, for each successive n, $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ should first be computed for arbitrary z_1, z_2, z_3, and z_4, with the factor $(z_1-z_3)(z_4-z_2)$ being explicitly canceled from both sides of (79), so that division by zero is avoided when boundary parameters need to be set to 1 in subsequent computations. Alternatively, expressions in which this cancelation has effectively been done, will be given in Section 4.2.

By replacing z_2 and z_4 by $\frac{1}{z_2}$ and $\frac{1}{z_4}$ respectively, Theorem 1 can be restated for the alternative quadruply-refined ASM generating function of (10) as

$$\begin{aligned}
&y(z_1-z_3)(z_2-z_4)\,\widetilde{Z}_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)\,Z_{n-2}(x,y)=\\
&\big((z_1-1)(z_2-1)+yz_1\big)\big((z_3-1)(z_4-1)+yz_3\big)\,(z_2z_4)^{n-1}\times\\
&\qquad\qquad Z_{n-1}^{\mathrm{adj}}(x,y;\tfrac{1}{z_4},z_1)\,Z_{n-1}^{\mathrm{adj}}(x,y;\tfrac{1}{z_2},z_3)-\\
&\big(x(z_4-1)(z_1-1)+yz_4\big)\big(x(z_2-1)(z_3-1)+yz_2\big)\,z_1z_3(z_2z_4)^{n-2}\times\\
&\qquad\qquad \widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,\tfrac{1}{z_2})\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_3,\tfrac{1}{z_4})-\\
&\quad(z_2-1)(z_3-1)\big((z_4-1)(z_1-1)+yz_1\big)\,(z_2z_4)^{n-1}\,Z_{n-1}^{\mathrm{adj}}(x,y;\tfrac{1}{z_4},z_1)\,Z_{n-2}(x,y)+\\
&\quad(z_3-1)(z_4-1)\big(x(z_1-1)(z_2-1)+yz_2\big)\,z_1z_2^{n-2}(xz_3)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,\tfrac{1}{z_2})\,Z_{n-2}(x,y)-\\
&\quad(z_4-1)(z_1-1)\big((z_2-1)(z_3-1)+yz_3\big)\,(z_2z_4)^{n-1}\,Z_{n-1}^{\mathrm{adj}}(x,y;\tfrac{1}{z_2},z_3)\,Z_{n-2}(x,y)+\\
&\quad(z_1-1)(z_2-1)\big(x(z_3-1)(z_4-1)+yz_4\big)\,z_3z_4^{n-2}(xz_1)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_3,\tfrac{1}{z_4})\,Z_{n-2}(x,y)+\\
&\qquad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\big((z_2z_4)^{n-1}-(x^2z_1z_3)^{n-1}\big)\,Z_{n-2}(x,y)^2. \qquad (80)
\end{aligned}$$

4.2. **Consequences for arbitrary bulk parameters x and y.**

In this section, some consequences of Theorem 1, for the case in which the bulk parameters x and y remain arbitrary, are derived. These results are obtained from (79) using only definitions and elementary properties of ASM generating functions from Section 2.2, and the initial conditions (from (8)–(10)) $Z_1^{\mathrm{adj}}(x,y;z_1,z_2)=\widetilde{Z}_1^{\mathrm{adj}}(x,y;z_1,z_2)=1$ and $Z_2^{\mathrm{adj}}(x,y;z_1,z_2)=\widetilde{Z}_2^{\mathrm{adj}}(x,y;z_1,z_2)=1+xz_1z_2$. In particular, most of the results are obtained by setting boundary parameters to 1, and using (11) for the corresponding specializations.

Alternative forms of some of the results of this section will be given in Section 5.7.

Corollary 1. *The triply-refined ASM generating function satisfies*

$$\begin{aligned}
&(z_2-z_1)(z_3-1)\,Z_n^{\mathrm{tri}}(x,y;z_1,z_2,z_3)\,Z_{n-2}(x,y)=\\
&\qquad\big((z_2-1)(z_3-1)-yz_2z_3\big)\,z_1\,Z_{n-1}^{\mathrm{adj}}(x,y;z_1,z_3)\,Z_{n-1}(x,y;z_2)-\\
&\qquad\big(x(z_1-1)(z_3-1)-y\big)\,z_1z_2z_3\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_2,z_3)\,Z_{n-1}(x,y;z_1)-\\
&\qquad\quad(z_1-1)(z_3-1)\,z_2\,Z_{n-1}(x,y;z_2)\,Z_{n-2}(x,y)+\\
&\qquad\qquad(z_2-1)(z_3-1)\,z_1\,(xz_2z_3)^{n-1}\,Z_{n-1}(x,y;z_1)\,Z_{n-2}(x,y). \qquad (81)
\end{aligned}$$

Proof. Set $z_2 = 1$ in (79) (and then relabel z_3 as z_2 and z_4 as z_3). □

Corollary 2. *The adjacent-boundary doubly-refined ASM generating functions satisfy the recursion relations*

$$\begin{aligned}(z_1-1)(z_2-1)\, Z_n^{\mathrm{adj}}(x,y;z_1,z_2)\, Z_{n-2}(x,y) &= yz_1z_2\, Z_{n-1}^{\mathrm{adj}}(x,y;z_1,z_2)\, Z_{n-1}(x,y) + \\ &\quad \big(x(z_1-1)(z_2-1)-y\big)\, z_1z_2\, Z_{n-1}(x,y;z_1)\, Z_{n-1}(x,y;z_2) + \\ &\quad (z_1-1)(z_2-1)\, Z_{n-1}(x,y)\, Z_{n-2}(x,y), \qquad (82)\\ (z_1-1)(z_2-1)\, \widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)\, Z_{n-2}(x,y) &= yz_1z_2\, \widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,z_2)\, Z_{n-1}(x,y) + \\ &\quad \big((z_1-1)(z_2-1)-yz_1z_2\big)\, Z_{n-1}(x,y;z_1)\, Z_{n-1}(x,y;z_2) + \\ &\quad (z_1-1)(z_2-1)\, (xz_1z_2)^{n-1}\, Z_{n-1}(x,y)\, Z_{n-2}(x,y). \qquad (83)\end{aligned}$$

Proof. To obtain (82), set $z_2 = 1$ in (81) (and then relabel z_3 as z_2). To obtain (83), set $z_1 = 1$ in (81) (and then relabel z_2 as z_1 and z_3 as z_2). □

Note that, if $Z_0(x,y)$ is taken to be 1, then (82) and (83) hold for all $n \geq 2$.

Corollary 3. *The triply-refined ASM generating function also satisfies*

$$\begin{aligned}y(z_2-z_1)z_3\, Z_n^{\mathrm{tri}}(x,y;z_1,z_2,z_3)\, Z_{n-1}(x,y) &= \\ &(z_1-1)\big((z_2-1)(z_3-1)-yz_2z_3\big)\, Z_n^{\mathrm{adj}}(x,y;z_1,z_3)\, Z_{n-1}(x,y;z_2) - \\ &(z_2-1)\big(x(z_1-1)(z_3-1)-y\big)\, z_1z_3\, \widetilde{Z}_n^{\mathrm{adj}}(x,y;z_2,z_3)\, Z_{n-1}(x,y;z_1) - \\ &(z_1-1)(z_2-1)(z_3-1)\, Z_{n-1}(x,y;z_2)\, Z_{n-1}(x,y) + \\ &(z_1-1)(z_2-1)(z_3-1)\, z_1z_2^{n-1}(xz_3)^n\, Z_{n-1}(x,y;z_1)\, Z_{n-1}(x,y). \qquad (84)\end{aligned}$$

Proof. Use (82)–(83) to replace $Z_{n-1}^{\mathrm{adj}}(x,y;z_1,z_3)$ and $\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_2,z_3)$ in (81) by terms which instead contain $Z_n^{\mathrm{adj}}(x,y;z_1,z_3)$ and $\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_2,z_3)$, and then cancel an overall factor which contains a term $z_3 - 1$. □

Corollary 4. *The opposite-boundary doubly-refined ASM generating function satisfies*

$$\begin{aligned}(z_1-z_2)\, Z_n^{\mathrm{opp}}(x,y;z_1,z_2)\, Z_{n-1}(x,y) &= (z_1-1)\, z_2\, Z_n(x,y;z_1)\, Z_{n-1}(x,y;z_2) - \\ &\quad z_1\, (z_2-1)\, Z_{n-1}(x,y;z_1)\, Z_n(x,y;z_2). \qquad (85)\end{aligned}$$

Proof. Set $z_3 = 1$ in (84). □

Note that Corollary 4 is a previously-known result, as already given in (61) and discussed in Section 3.4.

Corollary 5. *The adjacent-boundary doubly-refined ASM generating functions can be expressed as*

$$Z_n^{\mathrm{adj}}(x,y;z_1,z_2)=Z_{n-1}(x,y)\left(1+\sum_{i=1}^{n-1}\left(\frac{y\,z_1\,z_2}{(z_1-1)(z_2-1)}\right)^{n-i}\times\right.$$
$$\left.\left(1+\frac{(x(z_1-1)(z_2-1)-y)\,Z_i(x,y;z_1)\,Z_i(x,y;z_2)}{y\,Z_{i-1}(x,y)\,Z_i(x,y)}\right)\right),\quad(86)$$

$$\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)=Z_{n-1}(x,y)\left((xz_1z_2)^{n-1}+\sum_{i=1}^{n-1}\left(\frac{y}{(z_1-1)(z_2-1)}\right)^{n-i}\times\right.$$
$$\left.\left(x^{i-1}(z_1z_2)^{n-1}+\frac{(z_1z_2)^{n-i-1}((z_1-1)(z_2-1)-yz_1z_2)\,Z_i(x,y;z_1)\,Z_i(x,y;z_2)}{y\,Z_{i-1}(x,y)\,Z_i(x,y)}\right)\right),\quad(87)$$

where, in the sums over i, $Z_0(x,y)$ is taken to be 1.

Proof. It can be checked straightforwardly (again taking $Z_0(x,y)=1$), that the initial conditions $Z_1^{\mathrm{adj}}(x,y;z_1,z_2)=\widetilde{Z}_1^{\mathrm{adj}}(x,y;z_1,z_2)=1$ and recursion relations (82)–(83) are satisfied by (86)–(87). ☐

Corollary 6. *The two types of adjacent-boundary doubly-refined ASM generating function are related by*

$$((z_1-1)(z_2-1)-yz_1z_2)\,Z_n^{\mathrm{adj}}(x,y;z_1,z_2)-(x(z_1-1)(z_2-1)-y)\,z_1z_2\,\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)$$
$$=(z_1-1)(z_2-1)\big(1-(xz_1z_2)^n\big)\,Z_{n-1}(x,y).\quad(88)$$

Proof. This can be obtained directly from (86)–(87). Alternatively, it can be obtained from (82)–(83), by showing that $((z_1-1)(z_2-1)-yz_1z_2)\,Z_n^{\mathrm{adj}}(x,y;z_1,z_2)-(z_1-1)(z_2-1)\,Z_{n-1}(x,y)$ and $(x(z_1-1)(z_2-1)-y)\,z_1z_2\,\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)-(z_1-1)(z_2-1)(xz_1z_2)^n\,Z_{n-1}(x,y)$ satisfy the same recursion relation and initial condition. ☐

Note that, from (12), the two types of adjacent-boundary doubly-refined ASM generating function are also related by

$$Z_n^{\mathrm{adj}}(x,y;z_1,z_2)=x^{n(n-1)/2}\,(z_1z_2)^{n-1}\,\widetilde{Z}_n^{\mathrm{adj}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_1},\tfrac{1}{z_2}\big).\quad(89)$$

Combining (88) and (89), it follows that the adjacent-boundary doubly-refined ASM generating function satisfies

$$((z_1-1)(z_2-1)-yz_1z_2)\,Z_n^{\mathrm{adj}}(x,y;z_1,z_2)=$$
$$(x(z_1-1)(z_2-1)-y)\,(z_1z_2)^n\,x^{n(n-1)/2}\,Z_n^{\mathrm{adj}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_1},\tfrac{1}{z_2}\big)+$$
$$(z_1-1)(z_2-1)\big(1-(xz_1z_2)^n\big)\,Z_{n-1}(x,y).\quad(90)$$

By using (79) and (88), together with identities from (12), it is also possible to obtain a certain identity which involves $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ and $Z_n^{\mathrm{quad}}(x,y;z_3,z_2,z_1,z_4)$ (or $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ and $Z_n^{\mathrm{quad}}(x,y;z_1,z_4,z_3,z_2)$). A form of this identity will be given in (141).

Corollary 7. *The boundary parameter coefficients in the singly-refined ASM generating function, as defined in (14), satisfy*

$$\begin{aligned}Z_n(x,y)_k &= Z_{n-1}(x,y)\,\delta_{k,0}\,+\,Z_{n-1}(x,y)\sum_{i=0}^{k-1}\Biggl(y^{i+1}\binom{k-1}{i}\binom{n-1}{i+1}\,+\\&\qquad y^i\sum_{j_1=0}^{k-i-1}\,\sum_{j_2=0}^{n-i-2}\frac{Z_{n-i-1}(x,y)_{j_1}\,Z_{n-i-1}(x,y)_{j_2}}{Z_{n-i-1}(x,y)\,Z_{n-i-2}(x,y)}\Biggl(x\binom{k-j_1-2}{i-1}\binom{n-j_2-2}{i}\,-\\&\qquad\qquad y\binom{k-j_1-1}{i}\binom{n-j_2-1}{i+1}\Biggr)\Biggr),\qquad(91)\end{aligned}$$

where $Z_0(x,y)$, if it appears, is taken to be 1.

Proof. Expanding the factors $1/((z_1-1)(z_2-1))^{n-i}$ in (86) as binomial series, equating coefficients of $z_1^{k_1}z_2^{k_2}$ on both sides of (86), and using the definitions (14), gives

$$\begin{aligned}Z_n^{\mathrm{adj}}(x,y)_{k_1,k_2} &= Z_{n-1}(x,y)\,\delta_{k_1,0}\,\delta_{k_2,0}\,+\,Z_{n-1}(x,y)\sum_{i=0}^{\min(k_1,k_2)-1}\Biggl(y^{i+1}\binom{k_1-1}{i}\binom{k_2-1}{i}\,+\\&\qquad y^i\sum_{j_1=0}^{k_1-i-1}\,\sum_{j_2=0}^{k_2-i-1}\frac{Z_{n-i-1}(x,y)_{j_1}\,Z_{n-i-1}(x,y)_{j_2}}{Z_{n-i-1}(x,y)\,Z_{n-i-2}(x,y)}\Biggl(x\binom{k_1-j_1-2}{i-1}\binom{k_2-j_2-2}{i-1}\,-\\&\qquad\qquad y\binom{k_1-j_1-1}{i}\binom{k_2-j_2-1}{i}\Biggr)\Biggr).\qquad(92)\end{aligned}$$

Summing (92) over k_2, using a standard binomial coefficient summation identity, and relabeling k_1 as k, then gives (91). ☐

Corollary 8. *The unrefined ASM generating function satisfies*

$$\begin{aligned}Z_n(x,y) &= Z_{n-1}(x,y)\Biggl(1\,+\,\sum_{i=0}^{n-2}\Biggl(y^{i+1}\binom{n-1}{i+1}^2\,+\\&\qquad\frac{xy^i\bigl(\sum_{j=0}^{n-i-2}\binom{n-j-2}{i}Z_{n-i-1}(x,y)_j\bigr)^2-y^{i+1}\bigl(\sum_{j=0}^{n-i-2}\binom{n-j-1}{i+1}Z_{n-i-1}(x,y)_j\bigr)^2}{Z_{n-i-1}(x,y)\,Z_{n-i-2}(x,y)}\Biggr)\Biggr),\qquad(93)\end{aligned}$$

where $Z_0(x,y)$ is taken to be 1.

Proof. Sum (91) over k, and use a standard binomial coefficient summation identity. ☐

Note that various other expressions for $Z_n(x,y)_k$ and $Z_n(x,y)$, which provide alternatives to (91) and (93), can be obtained by taking the limits $z_1\to 1$ and $z_2\to 1$ of (86) in ways somewhat different from those used in Corollaries 7 and 8.

As observed in Section 4.1, the main result (79) enables all of the ASM generating functions of (7), (9) and (10) to be computed recursively. The corollaries of this section essentially comprise special cases of (79) which apply to specific ASM generating functions, and can alternatively be used for their computation.

For example, it can be seen that (91) and (93) give $Z_n(x,y)_k$ and $Z_n(x,y)$ in terms of $Z_i(x,y)_k$ and $Z_i(x,y)$ for $i=1,\dots,n-1$, and thereby enable the singly-refined and unrefined ASM generating functions to be computed recursively. The adjacent-boundary doubly-refined, opposite-boundary doubly-refined, triply-refined and quadruply-refined ASM generating functions can then be computed using (86)–(87), (85), (81) or (84), and (79), respectively.

Note that the opposite-boundary doubly-refined, singly-refined and unrefined ASM generating functions can instead be computed using the determinant formulae (53), (55) and (56).

4.3. **Consequences for bulk parameters $x=y=1$.** In this section, some consequences of Theorem 1, for the case in which the bulk parameters x and y are both set to 1, are derived. In contrast to the consequences of Theorem 1 given in Section 4.2 for the case of arbitrary x and y, which were obtained essentially using only (79), the consequences in this section are obtained using (79) together with the nontrivial relation (44) between the adjacent-boundary and opposite-boundary doubly-refined ASM generating functions at $x=y=1$.

Corollary 9. *The alternative quadruply-refined ASM generating function at $x=y=1$ satisfies*

$$\begin{aligned}
&(z_4z_1-z_4+1)(z_1z_2-z_1+1)(z_2z_3-z_2+1)(z_3z_4-z_3+1)\,\widetilde{Z}^{\mathrm{quad}}_n(1,1;z_1,z_2,z_3,z_4)=\\
&\frac{z_1\,z_2\,z_3\,z_4}{\mathcal{A}_{n-2}\,(z_1-z_3)(z_2-z_4)}\times\\
&\big((z_1z_2-z_1+1)(z_1z_2-z_2+1)(z_3z_4-z_3+1)(z_3z_4-z_4+1)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_4,z_1)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_2,z_3)-\\
&(z_4z_1-z_4+1)(z_4z_1-z_1+1)(z_2z_3-z_2+1)(z_2z_3-z_3+1)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_1,z_2)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_3,z_4)\big)+\\
&\qquad(z_2-1)(z_3-1)(z_1z_2-z_1+1)(z_3z_4-z_3+1)z_4z_1z_2^{n-1}Z^{\mathrm{opp}}_{n-1}(1,1;z_4,z_1)+\\
&\qquad(z_3-1)(z_4-1)(z_2z_3-z_2+1)(z_4z_1-z_4+1)z_1z_2z_3^{n-1}Z^{\mathrm{opp}}_{n-1}(1,1;z_1,z_2)+\\
&\qquad(z_4-1)(z_1-1)(z_3z_4-z_3+1)(z_1z_2-z_1+1)z_2z_3z_4^{n-1}Z^{\mathrm{opp}}_{n-1}(1,1;z_2,z_3)+\\
&\qquad(z_1-1)(z_2-1)(z_4z_1-z_4+1)(z_2z_3-z_2+1)z_3z_4z_1^{n-1}Z^{\mathrm{opp}}_{n-1}(1,1;z_3,z_4)+\\
&\quad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\big((z_1z_2-z_1+1)(z_3z_4-z_3+1)(z_2z_4)^{n-1}+\\
&\qquad\qquad(z_2z_3-z_2+1)(z_4z_1-z_4+1)(z_1z_3)^{n-1}\big)\mathcal{A}_{n-2}.
\end{aligned}\tag{94}$$

Proof. Setting $x=y=1$ in (80), and using (89), gives

$$\begin{aligned}
&(z_1-z_3)(z_2-z_4)\widetilde{Z}^{\mathrm{quad}}_n(1,1;z_1,z_2,z_3,z_4)\,\mathcal{A}_{n-2}=\\
&\qquad(z_1z_2-z_2+1)(z_3z_4-z_4+1)(z_2z_4)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_4},z_1)\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_2},z_3)-\\
&\qquad(z_4z_1-z_1+1)(z_2z_3-z_3+1)(z_1z_3)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_1},z_2)\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_3},z_4)-\\
&\qquad\quad(z_2-1)(z_3-1)(z_4z_1-z_4+1)(z_2z_4)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_4},z_1)\,\mathcal{A}_{n-2}+
\end{aligned}$$

$$\begin{aligned}&(z_3-1)(z_4-1)(z_1z_2-z_1+1)(z_1z_3)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_1},z_2)\,\mathcal{A}_{n-2}\,-\\&(z_4-1)(z_1-1)(z_2z_3-z_2+1)(z_2z_4)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_2},z_3)\,\mathcal{A}_{n-2}\,+\\&(z_1-1)(z_2-1)(z_3z_4-z_3+1)(z_1z_3)^{n-1}\,Z^{\mathrm{adj}}_{n-1}(1,1;\tfrac{1}{z_3},z_4)\,\mathcal{A}_{n-2}\,+\\&\qquad(z_1-1)(z_2-1)(z_3-1)(z_4-1)\big((z_2z_4)^{n-1}-(z_1z_3)^{n-1}\big)\,\mathcal{A}_{n-2}^2.\end{aligned}\tag{95}$$

Multiplying both sides of (95) by $(z_4z_1-z_4+1)(z_1z_2-z_1+1)(z_2z_3-z_2+1)(z_3z_4-z_3+1)/((z_1-z_3)(z_2-z_4)\mathcal{A}_{n-2})$, and using (44), in the form $(z_1z_2-z_1+1)z_1^{n-2}Z^{\mathrm{adj}}_{n-1}(1,1;\frac{1}{z_1},z_2)=z_2Z^{\mathrm{opp}}_{n-1}(1,1;z_1,z_2)+z_1^{n-2}(z_1-1)(z_2-1)\mathcal{A}_{n-2}$, then gives (94). ☐

Corollary 10. *The triply-refined ASM generating function at $x=y=1$ satisfies*

$$\begin{aligned}&(z_1z_3-z_3+1)(z_2z_3-z_2+1)z_3^{n-1}Z^{\mathrm{tri}}_n(1,1;z_1,z_2,\tfrac{1}{z_3})=\\&\quad\frac{z_1\,z_3}{\mathcal{A}_{n-2}\,(z_1-z_2)(z_3-1)}\Big((z_1z_3-z_1+1)(z_1z_3-z_3+1)z_2\,Z_{n-1}(1,1;z_1)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_2,z_3)\,-\\&\qquad\qquad(z_2z_3-z_2+1)(z_2z_3-z_3+1)z_1\,Z_{n-1}(1,1;z_2)\,Z^{\mathrm{opp}}_{n-1}(1,1;z_1,z_3)\Big)\,+\\&\qquad\qquad(z_2-1)(z_3-1)(z_1z_3-z_3+1)z_1z_2^{n-1}Z_{n-1}(1,1;z_1)\,+\\&\qquad\qquad\qquad(z_1-1)(z_3-1)(z_2z_3-z_2+1)z_3^{n-1}Z_{n-1}(1,1;z_2).\end{aligned}\tag{96}$$

Proof. Set $z_2=1$ in (94) (and then relabel z_3 as z_2 and z_4 as z_3). ☐

It can be seen that (94) and (96) provide alternative formulae to (48)–(50) for the quadruply- and triply-refined ASM generating functions at $x=y=1$. In fact, (94) and (96) differ from (50) and (49), respectively, only in the first terms on each RHS. It can also be seen that the last five terms on the RHS of (94) could be replaced by the last five terms on the RHS of (48).

5. Proofs

In this section, proofs of Theorem 1, and of several of the results of Section 3 are given. Preliminary results which are needed for these proofs are obtained or stated in Sections 5.1–5.5, while the main steps of the proofs are given in Sections 5.6 and 5.8–5.10. In Section 5.7, alternative forms of certain results of Section 4, and some further results, are discussed.

5.1. A generalized ASM generating function.

In this section, a generating function which generalizes the quadruply-refined ASM generating function is introduced, and some of its elementary properties are identified. This generating function will be shown, in Section 5.3, to be proportional to a certain case of the partition function of the six-vertex model with DWBC.

The generalized ASM generating function involves the six statistics of (3)–(4), together with four further statistics associated with the entries in the corners of an ASM, and is defined, for indeterminates x, y, z_1, z_2, z_3, z_4, z_{41}, z_{12}, z_{23} and z_{34}, as

$$Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) = \sum_{A\in\mathrm{ASM}(n)} x^{\nu(A)}\, y^{\mu(A)}\, z_1^{\rho_{\mathrm{T}}(A)}\, z_2^{\rho_{\mathrm{R}}(A)}\, z_3^{\rho_{\mathrm{B}}(A)}\, z_4^{\rho_{\mathrm{L}}(A)}\, {z_{41}}^{1-A_{11}}\, {z_{12}}^{1-A_{1n}}\, {z_{23}}^{1-A_{nn}}\, {z_{34}}^{1-A_{n1}}. \tag{97}$$

For example, for $n = 3$, the function is

$$Z_3^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) = z_{12}\,z_{34} + x\,z_1\,z_4\,z_{41}\,z_{12}\,z_{34} + x\,z_2\,z_3\,z_{12}\,z_{23}\,z_{34} + x^2\,z_1\,z_2\,z_3^2\,z_4^2\,z_{41}\,z_{12}\,z_{23} + x^2\,z_1^2\,z_2^2\,z_3\,z_4\,z_{41}\,z_{23}\,z_{34} + x^3\,z_1^2\,z_2^2\,z_3^2\,z_4^2\,z_{41}\,z_{23} + x\,y\,z_1\,z_2\,z_3\,z_4\,z_{41}\,z_{12}\,z_{23}\,z_{34}, \tag{98}$$

where the terms are written in an order which corresponds to that used in (2).

It can be seen immediately that the quadruply-refined ASM generating function is

$$Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4) = Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;1,1,1,1). \tag{99}$$

By acting on ASM(n) with transposition or anticlockwise quarter-turn rotation, and using (6), it follows that

$$\begin{aligned} Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) &= Z_n^{\mathrm{gen}}(x,y;z_4,z_3,z_2,z_1;z_{41},z_{34},z_{23},z_{12}) \\ &= x^{n(n-1)/2}\,(z_1z_2z_3z_4)^{n-1}\,Z_n^{\mathrm{gen}}\big(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_2},\tfrac{1}{z_3},\tfrac{1}{z_4},\tfrac{1}{z_1};z_{12},z_{23},z_{34},z_{41}\big). \end{aligned} \tag{100}$$

The properties of ASMs with a 1 in the top-left corner, as outlined in Section 2.1, imply that

$$\begin{aligned} Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;0,z_{12},z_{23},z_{34}) &= Z_n^{\mathrm{gen}}(x,y;0,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) \\ &= z_{12}\,z_{34}\,Z_{n-1}^{\mathrm{gen}}(x,y;1,z_2,z_3,1;1,1,z_{23},1), \end{aligned} \tag{101}$$

and the fact that an ASM cannot contain a 1 in both its top-left and top-right corner, implies that

$$Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;0,0,z_{23},z_{34}) = 0. \tag{102}$$

It can be seen from the definition (97) that $Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34})$ is linear in z_{41}, z_{12}, z_{23} and z_{34}. Therefore, using interpolation for these parameters at 0 and 1, it follows that

$$Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) = \sum_{i_1,i_2,i_3,i_4\in\{0,1\}} (1-z_{41})^{1-i_1}\,{z_{41}}^{i_1}\,(1-z_{12})^{1-i_2}\,{z_{12}}^{i_2}\,(1-z_{23})^{1-i_3}\,{z_{23}}^{i_3}\,(1-z_{34})^{1-i_4}\,{z_{34}}^{i_4} \times Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;i_1,i_2,i_3,i_3). \tag{103}$$

Applying cases of (9)–(10) and (99)–(102) to (103) (as a result of which, for example, (100) and (102) imply that nine of the sixteen terms on the RHS of (103) vanish, while (9) and (99)–(101) give $Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;0,1,0,1) = Z_{n-2}(x,y)$), it now follows that the generalized

ASM generating function can be expressed entirely in terms of quadruply-refined, adjacent-boundary doubly-refined and unrefined ASM generating functions as

$$\begin{aligned}Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34}) = {}& z_{41}\,z_{12}\,z_{23}\,z_{34}\,Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)+\\ & z_{41}\,z_{12}\,(1-z_{23})\,z_{34}\,Z_{n-1}^{\mathrm{adj}}(x,y;z_4,z_1)+\\ & z_{41}\,z_{12}\,z_{23}\,(1-z_{34})\,z_1z_2\,(xz_3z_4)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_1,z_2)+\\ & (1-z_{41})\,z_{12}\,z_{23}\,z_{34}\,Z_{n-1}^{\mathrm{adj}}(x,y;z_2,z_3)+\\ & z_{41}\,(1-z_{12})\,z_{23}\,z_{34}\,z_3z_4\,(xz_1z_2)^{n-1}\,\widetilde{Z}_{n-1}^{\mathrm{adj}}(x,y;z_3,z_4)+\\ & (1-z_{41})\,z_{12}\,(1-z_{23})\,z_{34}\,Z_{n-2}(x,y)+\\ & z_{41}\,(1-z_{12})\,z_{23}\,(1-z_{34})\,x^{2n-3}\,(z_1z_2z_3z_4)^{n-1}\,Z_{n-2}(x,y). \qquad (104)\end{aligned}$$

Some special cases of (104), which will be used in Section 5.6, are

$$\begin{aligned}Z_n^{\mathrm{gen}}(x,y;z_1,1,1,z_2;z_{12},1,1,1) &= Z_n^{\mathrm{gen}}(x,y;1,z_1,z_2,1;1,1,z_{12},1)\\ &= z_{12}\,Z_n^{\mathrm{adj}}(x,y;z_1,z_2)+(1-z_{12})\,Z_{n-1}(x,y),\\ Z_n^{\mathrm{gen}}(x,y;z_1,z_2,1,1;1,z_{12},1,1) &= Z_n^{\mathrm{gen}}(x,y;1,1,z_1,z_2;1,1,1,z_{12})\\ &= z_{12}\,\widetilde{Z}_n^{\mathrm{adj}}(x,y;z_1,z_2)+(1-z_{12})\,(xz_1z_2)^{n-1}\,Z_{n-1}(x,y). \qquad (105)\end{aligned}$$

5.2. **The bijection between ASMs and configurations of the six-vertex model with DWBC.** In this section, the set of configurations of the six-vertex model on an $n\times n$ grid with DWBC is described, and the details of a natural bijection between this set and ASM(n) are summarized. This is standard material, with similar accounts having been given, for example, by Behrend, Di Francesco and Zinn-Justin [16, Sec. 5.1], [17, Secs. 2.1 & 3.1].

The six-vertex, or square ice, model is a much-studied integrable statistical mechanical model (see, for example, Baxter [14, Chaps. 8 & 9] for further information and references), with DWBC for the model having been introduced and first studied by Korepin [85]. The bijection between ASMs and configurations of the model with DWBC was first discussed by Elkies, Kuperberg, Larsen and Propp [64, Sec. 7], with the details having mostly been observed previously, but using different terminology, by Robbins and Rumsey [125, pp. 179–180]. There exist closely-related bijections between ASMs and certain sets of osculating lattice paths (see Section 3.11 for references), and between ASMs and certain fully-packed loop configurations (see, for example, Propp [113, Sec. 7]).

Let $\mathcal{G}_n$ be the $n\times n$ undirected grid with vertex set $\{(i,j)\mid i,j=0,\ldots,n+1\}\setminus\{(0,0),(0,n+1),(n+1,0),(n+1,n+1)\}$, where (i,j) is taken to be in the ith row from the top and jth column from the left, and for which there are horizontal edges between (i,j) and $(i,j\pm1)$, and vertical edges between (i,j) and $(i\pm1,j)$, for each $i,j=1,\ldots n$. This grid is shown in Figure 1. Each vertex or edge of $\mathcal{G}_n$ can be described as either internal or external, i.e., the n^2 vertices of degree 4 are internal, the remaining $4n$ vertices of degree 1 are external, the $2n(n-1)$ edges which connect two internal vertices are internal, and the remaining $4n$ edges are external.



FIGURE 1. The grid $\mathcal{G}_n$.

A configuration of the six-vertex model on $\mathcal{G}_n$ with DWBC is an assignment of arrows to the edges of $\mathcal{G}_n$ such that the arrows on the external edges on the upper, right, lower and left boundaries of $\mathcal{G}_n$ are all directed upward, leftward, downward and rightward, respectively, while the arrows on the four edges incident to any internal vertex satisfy the condition that two point towards and two point away from the vertex.

Now define 6VDW(n) to be the set of all configurations of the six-vertex model on $\mathcal{G}_n$ with DWBC. For example,

6VDW(3) =



(106)

In an element of 6VDW(n), there are six possible configurations of arrows on the four edges incident to an internal vertex of $\mathcal{G}_n$. These so-called vertex configurations, are shown in Figure 2, where the numbers 1, . . . , 6 will be used to label the types, as indicated.



FIGURE 2. The possible arrow configurations on edges incident to an internal vertex.

For $C \in$ 6VDW(n), let $C_{ij} \in \{1,\dots,6\}$ denote the type of vertex configuration in C at internal vertex (i,j) of $\mathcal{G}_n$, and let $\mathcal{N}_{(k)}(C)$, $\mathcal{N}_{(k)}^{\mathrm{row}\,i}(C)$, $\mathcal{N}_{(k)}^{\mathrm{col}\,j}(C)$ and $\mathcal{N}_{(k)}^{ij}(C)$ denote the numbers of type-k vertex configurations in C in the whole of $\mathcal{G}_n$, in row i of $\mathcal{G}_n$, in column j of $\mathcal{G}_n$, or at vertex (i,j) of $\mathcal{G}_n$, respectively. Thus, for example, $\mathcal{N}_{(k)}^{ij}(C) = \delta_{C_{ij},k}$, and $\mathcal{N}_{(k)}(C) = |\{(i,j) \mid 1 \le i,j \le n,\ C_{ij} = k\}| = \sum_{i,j=1}^{n} \mathcal{N}_{(k)}^{ij}(C)$.

It can be shown easily that, for any $C \in 6\mathrm{VDW}(n)$,

$$\mathcal{N}_{(1)}(C)=\mathcal{N}_{(2)}(C), \quad \mathcal{N}_{(3)}(C)=\mathcal{N}_{(4)}(C), \quad \mathcal{N}_{(5)}(C)=\mathcal{N}_{(6)}(C)+n, \tag{107}$$

$$\mathcal{N}_{(5)}^{\mathrm{row}\,1}(C)=\mathcal{N}_{(5)}^{\mathrm{col}\,n}(C)=\mathcal{N}_{(5)}^{\mathrm{row}\,n}(C)=\mathcal{N}_{(5)}^{\mathrm{col}\,1}(C)=1. \tag{108}$$

(See, for example, Bressoud [22, p. 228], [23, p. 290] for a discussion of the first two equations of (107).) It can also be seen that, for any $C \in 6\mathrm{VDW}(n)$ and $1 \le i,j \le n$,

$$C_{1j} \in \{1,3,5\}, \quad C_{in} \in \{2,3,5\}, \quad C_{nj} \in \{2,4,5\}, \quad C_{i1} \in \{1,4,5\}, \tag{109}$$

and therefore that

$$C_{11} \in \{1,5\}, \quad C_{1n} \in \{3,5\}, \quad C_{nn} \in \{2,5\}, \quad C_{n1} \in \{4,5\}. \tag{110}$$

It can be shown straightforwardly that there is a natural bijection between $\mathrm{ASM}(n)$ and $6\mathrm{VDW}(n)$, and that, for each $A \in \mathrm{ASM}(n)$ and $C \in 6\mathrm{VDW}(n)$ which correspond under this bijection, the statistics (3)–(4) and (62) satisfy

$$\begin{gathered}\nu(A)=\mathcal{N}_{(1)}(C)\ (=\mathcal{N}_{(2)}(C)), \quad \mu(A)=\mathcal{N}_{(6)}(C)\ (=\mathcal{N}_{(5)}(C)-n),\\ \rho_{\mathrm{T}}(A)=\mathcal{N}_{(1)}^{\mathrm{row}\,1}(C), \ \ \rho_{\mathrm{R}}(A)=\mathcal{N}_{(2)}^{\mathrm{col}\,n}(C), \ \ \rho_{\mathrm{B}}(A)=\mathcal{N}_{(2)}^{\mathrm{row}\,n}(C), \ \ \rho_{\mathrm{L}}(A)=\mathcal{N}_{(1)}^{\mathrm{col}\,1}(C),\\ \nu^{\mathrm{row}\,i}(A)=\mathcal{N}_{(1)}^{\mathrm{row}\,i}(C)+\mathcal{N}_{(2)}^{\mathrm{row}\,i}(C), \quad \nu^{\mathrm{col}\,j}(A)=\mathcal{N}_{(1)}^{\mathrm{col}\,j}(C)+\mathcal{N}_{(2)}^{\mathrm{col}\,j}(C),\\ \mu^{\mathrm{row}\,i}(A)=\mathcal{N}_{(6)}^{\mathrm{row}\,i}(C), \quad \mu^{\mathrm{col}\,j}(A)=\mathcal{N}_{(6)}^{\mathrm{col}\,j}(C).\end{gathered} \tag{111}$$

The details of this bijection are as follows. To map $A \in \mathrm{ASM}(n)$ to $C \in 6\mathrm{VDW}(n)$, for each $i=1,\dots,n$ and $j=0,\dots,n$, assign a right or left arrow to the horizontal edge of $\mathcal{G}_n$ between (i,j) and $(i,j+1)$, according to whether the partial row sum $\sum_{j'=1}^{j} A_{ij'}$ is 0 or 1, respectively. Similarly, for each $i=0,\dots,n$ and $j=1,\dots,n$, assign an upward or downward arrow to the vertical edge of $\mathcal{G}_n$ between (i,j) and $(i+1,j)$, according to whether the partial column sum $\sum_{i'=1}^{i} A_{i'j}$ is 0 or 1, respectively. (The fact that each partial row sum and each column sum is 0 or 1 is given by (1).) To map $C \in 6\mathrm{VDW}(n)$ to $A \in \mathrm{ASM}(n)$, let

$$A_{ij}=\begin{cases}1, & C_{ij}=5\\ -1, & C_{ij}=6\\ 0, & C_{ij}\in\{1,2,3,4\}.\end{cases} \tag{112}$$

The behavior of the statistics μ, ρ_{T}, ρ_{R}, ρ_{B} and ρ_{R} under this bijection can be seen easily. To obtain the behavior of the statistics ν, $\nu^{\mathrm{row}\,i}$ and $\nu^{\mathrm{col}\,j}$, note that, for each $A \in \mathrm{ASM}(n)$ and $C \in 6\mathrm{VDW}(n)$ which correspond under the bijection, and each $1 \le i,j \le n$, $\sum_{i'=1}^{i-1} A_{i'j}=\sum_{j'=1}^{j} A_{ij'}=0$ (or equivalently $\sum_{i'=1}^{i} A_{i'j}=\sum_{j'=1}^{j-1} A_{ij'}=0$) if and only if $C_{ij}=1$, and $\sum_{i'=1}^{i-1} A_{i'j}=\sum_{j'=1}^{j} A_{ij'}=1$ (or equivalently $\sum_{i'=1}^{i} A_{i'j}=\sum_{j'=1}^{j-1} A_{ij'}=1$) if and only if $C_{ij}=2$.

As examples of this bijection, in (2) and (106) the elements of $\mathrm{ASM}(3)$ and $6\mathrm{VDW}(3)$ are listed in an order for which respective elements correspond under the bijection.

5.3. **The partition function of the six-vertex model with DWBC.** In this section, the partition function of the six-vertex model with DWBC is introduced. A relation between this partition function, for certain assignments of its parameters, and the generalized ASM generating function (97), for certain assignments of its parameters, is then derived using the bijection of Section 5.2.

Let a weight $W_{(k)}(u,v)$ be associated with the vertex configuration of type k, where u and v are so-called spectral parameters.

The partition function for the case of the six-vertex model of relevance here depends on these weights, and on spectral parameters u_i and v_j associated with row i and column j of $\mathcal{G}_n$, for each $1 \le i,j \le n$. Specifically, this partition function is defined as

$$Z_n^{\mathrm{6V}}(u_1,\dots,u_n;v_1,\dots,v_n) = \sum_{C\in\mathrm{6VDW}(n)} \prod_{i,j=1}^{n} W_{(C_{ij})}(u_i,v_j). \tag{113}$$

Let the weights now satisfy

$$\begin{gathered} W_{(1)}(u,v)=W_{(2)}(u,v)=a(u,v), \quad W_{(3)}(u,v)=W_{(4)}(u,v)=b(u,v), \\ W_{(5)}(u,v)=W_{(6)}(u,v)=c(u,v), \end{gathered} \tag{114}$$

for functions a, b and c.

Taking the spectral parameters in (113) to be

$$u_2=\ldots=u_{n-1}=r, \quad v_2=\ldots=v_{n-1}=s, \quad u_1=t_1, \quad v_n=t_2, \quad u_n=t_3, \quad v_1=t_4, \tag{115}$$

for indeterminates r, s and $t_1,\dots,t_4$, and using (107)–(110), gives

$$\begin{aligned} Z_n^{\mathrm{6V}}(t_1,r,\dots,r,t_3;t_4,s,\dots,s,t_2) = \sum_{C\in\mathrm{6VDW}(n)} a(r,s)^{2\mathcal{N}_{(1)}(C)}\, b(r,s)^{2\mathcal{N}_{(3)}(C)}\, c(r,s)^{2\mathcal{N}_{(6)}(C)+n} \times \\ \Big(\tfrac{a(t_1,s)}{a(r,s)}\Big)^{\mathcal{N}_{(1)}^{\mathrm{row}\,1}(C)-\mathcal{N}_{(1)}^{11}(C)} \Big(\tfrac{b(t_1,s)}{b(r,s)}\Big)^{\mathcal{N}_{(3)}^{\mathrm{row}\,1}(C)-\mathcal{N}_{(3)}^{1n}(C)} \Big(\tfrac{c(t_1,s)}{c(r,s)}\Big)^{1-\mathcal{N}_{(5)}^{11}(C)-\mathcal{N}_{(5)}^{1n}(C)} \times \\ \Big(\tfrac{a(r,t_2)}{a(r,s)}\Big)^{\mathcal{N}_{(2)}^{\mathrm{col}\,n}(C)-\mathcal{N}_{(2)}^{nn}(C)} \Big(\tfrac{b(r,t_2)}{b(r,s)}\Big)^{\mathcal{N}_{(3)}^{\mathrm{col}\,n}(C)-\mathcal{N}_{(3)}^{1n}(C)} \Big(\tfrac{c(r,t_2)}{c(r,s)}\Big)^{1-\mathcal{N}_{(5)}^{1n}(C)-\mathcal{N}_{(5)}^{nn}(C)} \times \\ \Big(\tfrac{a(t_3,s)}{a(r,s)}\Big)^{\mathcal{N}_{(2)}^{\mathrm{row}\,n}(C)-\mathcal{N}_{(2)}^{nn}(C)} \Big(\tfrac{b(t_3,s)}{b(r,s)}\Big)^{\mathcal{N}_{(4)}^{\mathrm{row}\,n}(C)-\mathcal{N}_{(4)}^{n1}(C)} \Big(\tfrac{c(t_3,s)}{c(r,s)}\Big)^{1-\mathcal{N}_{(5)}^{n1}(C)-\mathcal{N}_{(5)}^{nn}(C)} \times \\ \Big(\tfrac{a(r,t_4)}{a(r,s)}\Big)^{\mathcal{N}_{(1)}^{\mathrm{col}\,1}(C)-\mathcal{N}_{(1)}^{11}(C)} \Big(\tfrac{b(r,t_4)}{b(r,s)}\Big)^{\mathcal{N}_{(4)}^{\mathrm{col}\,1}(C)-\mathcal{N}_{(4)}^{n1}(C)} \Big(\tfrac{c(r,t_4)}{c(r,s)}\Big)^{1-\mathcal{N}_{(5)}^{11}(C)-\mathcal{N}_{(5)}^{n1}(C)} \times \\ \Big(\tfrac{a(t_1,t_4)}{a(r,s)}\Big)^{\mathcal{N}_{(1)}^{11}(C)} \Big(\tfrac{c(t_1,t_4)}{c(r,s)}\Big)^{\mathcal{N}_{(5)}^{11}(C)} \Big(\tfrac{b(t_1,t_2)}{b(r,s)}\Big)^{\mathcal{N}_{(3)}^{1n}(C)} \Big(\tfrac{c(t_1,t_2)}{c(r,s)}\Big)^{\mathcal{N}_{(5)}^{1n}(C)} \times \\ \Big(\tfrac{a(t_3,t_2)}{a(r,s)}\Big)^{\mathcal{N}_{(2)}^{nn}(C)} \Big(\tfrac{c(t_3,t_2)}{c(r,s)}\Big)^{\mathcal{N}_{(5)}^{nn}(C)} \Big(\tfrac{b(t_3,t_4)}{b(r,s)}\Big)^{\mathcal{N}_{(4)}^{n1}(C)} \Big(\tfrac{c(t_3,t_4)}{c(r,s)}\Big)^{\mathcal{N}_{(5)}^{n1}(C)}. \end{aligned} \tag{116}$$

It can be seen that

$$\mathcal{N}_{(1)}(C)+\mathcal{N}_{(3)}(C)+\mathcal{N}_{(6)}(C)=\tfrac{n(n-1)}{2}, \tag{117}$$

$$\begin{multline}\mathcal{N}^{\mathrm{row}\,1}_{(1)}(C)+\mathcal{N}^{\mathrm{row}\,1}_{(3)}(C)=\mathcal{N}^{\mathrm{col}\,n}_{(2)}(C)+\mathcal{N}^{\mathrm{col}\,n}_{(3)}(C)=\\ \mathcal{N}^{\mathrm{row}\,n}_{(2)}(C)+\mathcal{N}^{\mathrm{row}\,n}_{(4)}(C)=\mathcal{N}^{\mathrm{col}\,1}_{(1)}(C)+\mathcal{N}^{\mathrm{col}\,1}_{(4)}(C)=n-1, \tag{118}\end{multline}$$

$$\begin{multline}\mathcal{N}^{11}_{(1)}(C)+\mathcal{N}^{11}_{(5)}(C)=\mathcal{N}^{1n}_{(3)}(C)+\mathcal{N}^{1n}_{(5)}(C)=\\ \mathcal{N}^{nn}_{(2)}(C)+\mathcal{N}^{nn}_{(5)}(C)=\mathcal{N}^{n1}_{(4)}(C)+\mathcal{N}^{n1}_{(5)}(C)=1, \tag{119}\end{multline}$$

where (117) follows from (107) and $\sum_{k=1}^{6}\mathcal{N}_{(k)}(C)=n^2$, (118) follows from (108)–(109) and $\sum_{k=1}^{6}\mathcal{N}^{\mathrm{row}\,i}_{(k)}(C)=\sum_{k=1}^{6}\mathcal{N}^{\mathrm{col}\,j}_{(k)}(C)=n$, and (119) follows from (110).

By using (117)–(119) to eliminate $\mathcal{N}_{(3)}(C)$, $\mathcal{N}^{\mathrm{row}\,1}_{(3)}(C)$, $\mathcal{N}^{\mathrm{col}\,n}_{(3)}(C)$, $\mathcal{N}^{\mathrm{row}\,n}_{(4)}(C)$, $\mathcal{N}^{\mathrm{col}\,1}_{(4)}(C)$, $\mathcal{N}^{11}_{(5)}(C)$, $\mathcal{N}^{1n}_{(5)}(C)$, $\mathcal{N}^{nn}_{(5)}(C)$ and $\mathcal{N}^{n1}_{(5)}(C)$ from (116), and then using the bijection of Section 5.2 between $\mathrm{ASM}(n)$ and $\mathrm{6VDW}(n)$, the behavior (111) of the statistics (3)–(4) under this bijection, and the definition (97) of the generalized ASM generating function, it now follows that (116) can be written as

$$\begin{multline}Z^{\mathrm{6V}}_n(t_1,r,\ldots,r,t_3;t_4,s,\ldots,s,t_2)=\\ b(r,s)^{n(n-1)}\,c(r,s)^n\left(\tfrac{b(t_1,s)\,b(r,t_2)\,b(t_3,s)\,b(r,t_4)}{b(r,s)^4}\right)^{n-1}\tfrac{c(t_1,t_4)\,c(t_1,t_2)\,c(t_3,t_2)\,c(t_3,t_4)}{c(t_1,s)\,c(r,t_2)\,c(t_3,s)\,c(r,t_4)}\times\\ Z^{\mathrm{gen}}_n\Bigl(\left(\tfrac{a(r,s)}{b(r,s)}\right)^2,\left(\tfrac{c(r,s)}{b(r,s)}\right)^2;\ \tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\ \tfrac{a(r,t_2)\,b(r,s)}{a(r,s)\,b(r,t_2)},\ \tfrac{a(t_3,s)\,b(r,s)}{a(r,s)\,b(t_3,s)},\ \tfrac{a(r,t_4)\,b(r,s)}{a(r,s)\,b(r,t_4)};\ \tfrac{a(r,s)\,a(t_1,t_4)\,c(t_1,s)\,c(r,t_4)}{a(t_1,s)\,a(r,t_4)\,c(r,s)\,c(t_1,t_4)},\\ \tfrac{b(r,s)\,b(t_1,t_2)\,c(t_1,s)\,c(r,t_2)}{b(t_1,s)\,b(r,t_2)\,c(r,s)\,c(t_1,t_2)},\ \tfrac{a(r,s)\,a(t_3,t_2)\,c(t_3,s)\,c(r,t_2)}{a(t_3,s)\,a(r,t_2)\,c(r,s)\,c(t_3,t_2)},\ \tfrac{b(r,s)\,b(t_3,t_4)\,c(t_3,s)\,c(r,t_4)}{b(t_3,s)\,b(r,t_4)\,c(r,s)\,c(t_3,t_4)}\Bigr). \tag{120}\end{multline}$$

5.4. **The Izergin–Korepin determinant formula.** In this section, the Izergin–Korepin formula for the partition function (113), with certain assignments of the weights (114), is stated.

It was shown by Izergin [81], using certain results of Korepin [85], that if the weights (114) satisfy the Yang–Baxter equation (see, for example, Baxter [14, pp. 187–189]), then the partition function (113) is given by an explicit formula involving the determinant of a certain $n\times n$ matrix.

Let the functions a, b and c in (114) be given by

$$\begin{aligned}a(u,v)&=u\,q^{1/2}-v\,q^{-1/2},\\ b(u,v)&=v\,q^{1/2}-u\,q^{-1/2}=a(v,u),\\ c(u,v)&=\bigl(q-q^{-1}\bigr)\,u^{1/2}\,v^{1/2},\end{aligned} \tag{121}$$

where q is a further indeterminate, often known as the crossing parameter. (Note that q will be present in many subsequent equations, even though it may not appear explicitly.)

It can be seen that these functions satisfy

$$\frac{a(u,v)^2+b(u,v)^2-c(u,v)^2}{a(u,v)\,b(u,v)}=-(q+q^{-1}). \tag{122}$$

The fact that the LHS of (122) (which is often denoted as 2Δ) is independent of u and v implies that the six-vertex model weights given by (114) and (121) satisfy the Yang–Baxter equation (see, for example, Baxter [14, Eq. (9.6.14)]).

The resulting Izergin–Korepin determinant formula is then given by the following result.

Theorem (Izergin)**.** *The partition function* (113)*, with weights given by* (114) *and* (121)*, satisfies*

$$Z_n^{\text{6V}}(u_1,\ldots,u_n;v_1,\ldots,v_n)=\frac{\prod_{i,j=1}^{n}a(u_i,v_j)\,b(u_i,v_j)}{\prod_{1\le i<j\le n}(u_i-u_j)(v_j-v_i)}\det_{1\le i,j\le n}\left(\frac{c(u_i,v_j)}{a(u_i,v_j)\,b(u_i,v_j)}\right). \tag{123}$$

This theorem can be proved by showing that each side of (123) satisfies, and is uniquely determined by, four particular properties. Specifically, each side is symmetric in $u_1,\ldots,u_n$ and in $v_1,\ldots,v_n$ (which can be obtained for the LHS using the Yang–Baxter equation, and is immediate for the RHS), is a polynomial of degree $2n-1$ in each of $u_1^{1/2},\ldots,u_n^{1/2},v_1^{1/2},\ldots,v_n^{1/2}$, satisfies the same recursion relation for cases in which $u_i=q^{\pm1}v_j$ for some i and j, and satisfies the same initial condition at $n=1$. For further details of this proof, see, for example, Izergin [81], Izergin, Coker and Korepin [82, Sec. 5], Fonseca and Zinn-Justin [75, Sec. B.1], Kuperberg [92, Sec. 1], or Zinn-Justin [149, Secs. 2.5.2–2.5.3]. For an alternative proof, see Bogoliubov, Pronko and Zvonarev [20, Sec. 4].

Note that, although the determinant and the denominator of the prefactor on the RHS of (123) both vanish if $u_i=u_j$ or $v_i=v_j$ for some $i\neq j$, the RHS has a well-defined limit in these cases, as a polynomial in $u_1^{1/2},\ldots,u_n^{1/2},v_1^{1/2},\ldots,v_n^{1/2}$. Accordingly, it will be valid to use (123) to derive properties of the partition function for the assignments (115), as will be done in Section 5.6.

5.5. The Desnanot–Jacobi identity.

In this section, the Desnanot–Jacobi determinant identity is stated, and discussed briefly.

For a square matrix M, let M_{TL}, M_{TR}, M_{BR}, M_{BL} and M_{C} be the submatrices of M defined in Section 3.2. The Desnanot–Jacobi identity, in the form which will be used here, can be stated as follows.

Theorem (Desnanot–Jacobi)**.** *For any square matrix M,*

$$\det M\ \det M_{\text{C}}=\det M_{\text{TL}}\ \det M_{\text{BR}}-\det M_{\text{TR}}\ \det M_{\text{BL}}. \tag{124}$$

For an algebraic proof of (124), see, for example, Bressoud [22, Sec. 3.5], and for a combinatorial proof, see Zeilberger [147]. Cases of (124) for $n\times n$ matrices M with small values of n were published by Desnanot in 1819 (see Muir [108, Eqs. (A)–(G), (A′)–(G′), pp. 139–142]). The further attribution to Jacobi is based on the fact that, for arbitrary n, (124) corresponds to the case $m=2$ of the identity, published by Jacobi in 1834 (see Muir [108, Eq. (XX. 4), p. 208]), that for any $n\times n$ matrix M and any $m\le n$, each $m\times m$ minor of the matrix of $(n-1)\times(n-1)$ minors of M equals the complementary minor of M multiplied by $(\det M)^{m-1}$. For proofs of the Jacobi identity, using a variety of methods, see, for example, Brualdi and Schneider [27, Sec. 4], Knuth [84, Eq. (3.16)], Leclerc [100, Sec. 3.2], Muir [109, Sec. 175], or Turnbull [143, pp. 77–79].

For a discussion of alternative forms of (124), and of further determinant identities of which (124) is a special case, see, for example, Behrend, Di Francesco and Zinn-Justin [16, Sec. 4].

5.6. **Proof of Theorem 1.** In this section, the main steps in the proof of Theorem 1 are given. These involve using the relation (120) between the generalized ASM generating function and the partition function of the six-vertex model with DWBC, the Izergin–Korepin formula (123), and the Desnanot–Jacobi identity (124).

By applying the Desnanot–Jacobi identity (124) to the matrix $\left(\frac{c(u_i,v_j)}{a(u_i,v_j)\,b(u_i,v_j)}\right)_{1\le i,j\le n}$, and then applying the Izergin–Korepin formula (123) to each of the six determinants which appear, it follows that the partition function (113), with weights given by (114) and (121), satisfies

$$\begin{aligned}(u_1-u_n)\,(v_n-v_1)\,Z^{6\mathrm{V}}_n(u_1\ldots,u_n;v_1,\ldots,v_n)\,Z^{6\mathrm{V}}_{n-2}(u_2,\ldots,u_{n-1};v_2,\ldots,v_{n-1}) ={}&\\ a(u_1,v_n)\,b(u_1,v_n)\,a(u_n,v_1)\,b(u_n,v_1)\,Z^{6\mathrm{V}}_{n-1}(u_1,\ldots,u_{n-1};v_1,\ldots,v_{n-1})\times&\\ Z^{6\mathrm{V}}_{n-1}(u_2,\ldots,u_n;v_2,\ldots,v_n)&\\ -\;a(u_1,v_1)\,b(u_1,v_1)\,a(u_n,v_n)\,b(u_n,v_n)\,Z^{6\mathrm{V}}_{n-1}(u_1,\ldots,u_{n-1};v_2,\ldots,v_n)\times&\\ Z^{6\mathrm{V}}_{n-1}(u_2,\ldots,u_n;v_1,\ldots,v_{n-1}),&\end{aligned}\tag{125}$$

for any $u_1,\ldots,u_n,v_1,\ldots,v_n$. In fact, it can be seen, from the previous derivation, that (125) is satisfied by any function which has the form of (123), for arbitrary functions a, b and c.

Certain forms of the Desnanot–Jacobi identity have previously been combined with certain cases of the Izergin–Korepin formula by Korepin and Zinn-Justin [86, Sec. 3] (see also Sogo [129, Sec. 4]), and by Behrend, Di Francesco and Zinn-Justin [16, Sec. 5.4].

Now let parameters x, y, z_1, z_2, z_3, z_4, z_{41}, z_{12}, z_{23}, z_{34}, z'_{41}, z'_{12}, z'_{23} and z'_{34} be given in terms of parameters q, r, s, t_1, t_2, t_3 and t_4 by

$$\begin{gathered}x=\left(\tfrac{a(r,s)}{b(r,s)}\right)^2,\qquad y=\left(\tfrac{c(r,s)}{b(r,s)}\right)^2,\\ z_1=\tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\quad z_2=\tfrac{a(r,t_2)\,b(r,s)}{a(r,s)\,b(r,t_2)},\quad z_3=\tfrac{a(t_3,s)\,b(r,s)}{a(r,s)\,b(t_3,s)},\quad z_4=\tfrac{a(r,t_4)\,b(r,s)}{a(r,s)\,b(r,t_4)},\\ z_{41}=\tfrac{a(r,s)\,a(t_1,t_4)}{a(t_1,s)\,a(r,t_4)},\quad z_{12}=\tfrac{b(r,s)\,b(t_1,t_2)}{b(t_1,s)\,b(r,t_2)},\quad z_{23}=\tfrac{a(r,s)\,a(t_3,t_2)}{a(t_3,s)\,a(r,t_2)},\quad z_{34}=\tfrac{b(r,s)\,b(t_3,t_4)}{b(t_3,s)\,b(r,t_4)},\\ z'_{41}=\tfrac{b(r,s)\,b(t_1,t_4)}{b(t_1,s)\,b(r,t_4)},\quad z'_{12}=\tfrac{a(r,s)\,a(t_1,t_2)}{a(t_1,s)\,a(r,t_2)},\quad z'_{23}=\tfrac{b(r,s)\,b(t_3,t_2)}{b(t_3,s)\,b(r,t_2)},\quad z'_{34}=\tfrac{a(r,s)\,a(t_3,t_4)}{a(t_3,s)\,a(r,t_4)},\end{gathered}\tag{126}$$

where the functions a, b and c are given by (121).

The parameterizations (126) allow arbitrary x, y and $z_1,\ldots,z_4$ to be expressed in terms of q, r, s and $t_1,\ldots,t_4$. In particular, it follows, using (121), (122) and (126), that q can first be taken to be a solution of

$$x^{1/2}\,q^2+(x-y+1)\,q+x^{1/2}=0,\tag{127}$$

that r and s can then be taken to satisfy

$$(x^{1/2}+q)\,r=(x^{1/2}\,q+1)\,s, \tag{128}$$

and that $t_1,\ldots,t_4$ are then given by

$$t_1=\tfrac{s(x^{1/2}\,z_1\,q+1)}{x^{1/2}\,z_1+q},\quad t_2=\tfrac{r(x^{1/2}\,z_2+q)}{x^{1/2}\,z_2\,q+1},\quad t_3=\tfrac{s(x^{1/2}\,z_3\,q+1)}{x^{1/2}\,z_3+q},\quad t_4=\tfrac{r(x^{1/2}\,z_4+q)}{x^{1/2}\,z_4\,q+1}. \tag{129}$$

It can be checked easily that the parameterizations (126) imply that z_{41}, z_{12}, z_{23}, z_{34}, z'_{41}, z'_{12}, z'_{23} and z'_{34} can be expressed in terms of x, y and $z_1,\ldots,z_4$ as

$$z_{41}=1-\tfrac{(z_4-1)(z_1-1)}{y\,z_4\,z_1},\quad z_{12}=1-\tfrac{x(z_1-1)(z_2-1)}{y},\quad z_{23}=1-\tfrac{(z_2-1)(z_3-1)}{y\,z_2\,z_3},\quad z_{34}=1-\tfrac{x(z_3-1)(z_4-1)}{y}, \tag{130}$$

and

$$z'_{41}=1-\tfrac{x(z_4-1)(z_1-1)}{y},\quad z'_{12}=1-\tfrac{(z_1-1)(z_2-1)}{y\,z_1\,z_2},\quad z'_{23}=1-\tfrac{x(z_2-1)(z_3-1)}{y},\quad z'_{34}=1-\tfrac{(z_3-1)(z_4-1)}{y\,z_3\,z_4}. \tag{131}$$

Thus, each z_{ij} and z'_{ij} has the form $f(y,z_i,z_j)=1-\frac{(z_i-1)(z_j-1)}{y\,z_i\,z_j}$ or $f(\frac{y}{x},\frac{1}{z_i},\frac{1}{z_j})$.

The parameterizations (126) also give

$$(t_2-t_4)\,(t_1-t_3)=\tfrac{a(r,s)^2\,b(t_1,s)\,b(r,t_2)\,b(t_3,s)\,b(r,t_4)}{(b(r,s)\,c(r,s))^2}\,(z_4-z_2)\,(z_1-z_3). \tag{132}$$

Assigning the parameters in (125) according to (115), applying the relation (120) to each of the six cases of the partition function in (125), and using (126) and (132), it now follows that

$$\begin{aligned}(z_4-z_2)\,(z_1-z_3)\,Z^{\mathrm{gen}}_n(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34})\,Z^{\mathrm{gen}}_{n-2}(x,y;1,1,1,1;1,1,1,1)=\\ y\,z_1\,z_2\,z_3\,z_4\,\bigl(z_{12}\,z'_{12}\,z_{34}\,z'_{34}\,Z^{\mathrm{gen}}_{n-1}(x,y;z_1,1,1,z_4;z_{41},1,1,1)\times\\ Z^{\mathrm{gen}}_{n-1}(x,y;1,z_2,z_3,1;1,1,z_{23},1)\\ -\;z_{41}\,z'_{41}\,z_{23}\,z'_{23}\,Z^{\mathrm{gen}}_{n-1}(x,y;z_1,z_2,1,1;1,z_{12},1,1)\times\\ Z^{\mathrm{gen}}_{n-1}(x,y;1,1,z_3,z_4;1,1,1,z_{34})\bigr),\end{aligned} \tag{133}$$

where x, y and $z_1,\ldots,z_4$ are arbitrary, while the remaining parameters are given by (130)–(131).

Finally, the required identity (79) follows from (133) by applying (104), including the special cases (105), to each of the six cases of generalized ASM generating functions, and using (130)–(131) to eliminate z_{41}, z_{12}, z_{23}, z_{34}, z'_{41}, z'_{12}, z'_{23} and z'_{34}.

5.7. **Alternative statement of some results of Section 4.1.** In this section, it is shown that some of the results of Section 4.1 can be stated, slightly more compactly, in terms of a certain function which will be defined in (134) (and has essentially already appeared in 133). Also, an additional symmetry property for ASM generating functions is obtained, and further quadratic relations satisfied by these functions are discussed briefly.

Define a function

$$\begin{aligned}Y_n(x,y;z_1,z_2,z_3,z_4)=\sum_{A\in\mathrm{ASM}(n)} & x^{\nu(A)}\,y^{\mu(A)+A_{11}+A_{1n}+A_{nn}+A_{n1}}\times\\ & z_1^{\rho_{\mathrm{T}}(A)+A_{11}-1}\,z_2^{\rho_{\mathrm{R}}(A)+A_{nn}-1}\,z_3^{\rho_{\mathrm{B}}(A)+A_{nn}-1}\,z_4^{\rho_{\mathrm{L}}(A)+A_{11}-1}\times\\ & \quad(yz_4z_1-(z_4-1)(z_1-1))^{1-A_{11}}\,(y-x(z_1-1)(z_2-1))^{1-A_{1n}}\times\\ & \qquad(yz_2z_3-(z_2-1)(z_3-1))^{1-A_{nn}}\,(y-x(z_3-1)(z_4-1))^{1-A_{n1}}.\end{aligned}\tag{134}$$

It follows that $Y_n(x,y;z_1,z_2,z_3,z_4)$ is a polynomial in x, y and $z_1,\ldots,z_4$, with integer coefficients.

Also define

$$Y_n(x,y;z_1,z_2)=Y_n(x,y;z_1,z_2,1,1).\tag{135}$$

It can be seen, using the definition (97) of the generalized ASM generating function, that

$$\begin{aligned}Y_n(x,y;z_1,z_2,z_3,z_4)=y^4Z_n^{\mathrm{gen}}\bigl(&x,y;z_1,z_2,z_3,z_4;\\ &1-\tfrac{(z_4-1)(z_1-1)}{y\,z_4\,z_1},1-\tfrac{x(z_1-1)(z_2-1)}{y},1-\tfrac{(z_2-1)(z_3-1)}{y\,z_2\,z_3},1-\tfrac{x(z_3-1)(z_4-1)}{y}\bigr),\end{aligned}\tag{136}$$

i.e., $Y_n(x,y;z_1,z_2,z_3,z_4)=y^4Z_n^{\mathrm{gen}}(x,y;z_1,z_2,z_3,z_4;z_{41},z_{12},z_{23},z_{34})$, with z_{41}, z_{12}, z_{23} and z_{34} given by (130).

Therefore, by applying (104) to the RHS of (136), $Y_n(x,y;z_1,z_2,z_3,z_4)$ can be expressed in terms of quadruply-refined, adjacent-boundary doubly-refined and unrefined ASM generating functions.

It can be seen immediately from (134)–(135) that

$$\begin{aligned}Y_n(x,y;z_1,1,z_2,1)&=y^4\,Z_n^{\mathrm{opp}}(x,y;z_1,z_2),\\ Y_n(x,y;z,1)&=y^4\,Z_n(x,y;z),\\ Y_n(x,y;1,1)&=y^4\,Z_n(x,y),\end{aligned}\tag{137}$$

and by acting on ASM(n) in (134) with transposition or anticlockwise quarter-turn rotation and using (6) (or by applying (100) to (136)), it follows that

$$\begin{aligned}Y_n(x,y;z_1,z_2,z_3,z_4)&=Y_n(x,y;z_4,z_3,z_2,z_1)\\ &=x^{n(n-1)/2+4}\,(z_1z_2z_3z_4)^{n-1}\,Y_n\bigl(\tfrac{1}{x},\tfrac{y}{x};\tfrac{1}{z_2},\tfrac{1}{z_3},\tfrac{1}{z_4},\tfrac{1}{z_1}\bigr).\end{aligned}\tag{138}$$

It can also be shown, using (134) and the properties of ASMs in which a 1 on a boundary is in a corner or separated from a corner by a single zero (as discussed in Section 2.1), that

$$Y_n(x,y;z_1,z_2,z_3,0)=\bigl(1+\tfrac{x(z_1-1)}{y}\bigr)\,\bigl(1+\tfrac{x(z_3-1)}{y}\bigr)\,Y_{n-1}(x,y;z_1,z_2,z_3,1).\tag{139}$$

Taking x, y, $z_1,\dots,z_4$, z_{41}, z_{12}, z_{23} and z_{34} to be parameterized by (126), and using (120), (130) and (136), gives

$$\begin{multline}
b(r,s)^{(n-1)(n-4)+8}\,c(r,s)^{n-8}\,\bigl(b(t_1,s)\,b(r,t_2)\,b(t_3,s)\,b(r,t_4)\bigr)^{n-1}\,t_1^{1/2}\,t_2^{1/2}\,t_3^{1/2}\,t_4^{1/2}\,(rs)^{-1}\times\\
Y_n\Bigl(\bigl(\tfrac{a(r,s)}{b(r,s)}\bigr)^2,\bigl(\tfrac{c(r,s)}{b(r,s)}\bigr)^2;\tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\tfrac{a(r,t_2)\,b(r,s)}{a(r,s)\,b(r,t_2)},\tfrac{a(t_3,s)\,b(r,s)}{a(r,s)\,b(t_3,s)},\tfrac{a(r,t_4)\,b(r,s)}{a(r,s)\,b(r,t_4)}\Bigr)=\\
Z_n^{\mathrm{6V}}(t_1,r,\dots,r,t_3;t_4,s,\dots,s,t_2),\quad(140)
\end{multline}$$

where the weights in the partition function are given by (114), and the functions a, b and c are given by (121).

It now follows from the symmetry of the partition function on the RHS of (140) in t_1 and t_3, and in t_2 and t_4, as discussed after (123), that

$$Y_n(x,y;z_1,z_2,z_3,z_4)=Y_n(x,y;z_3,z_2,z_1,z_4)=Y_n(x,y;z_1,z_4,z_3,z_2).\qquad(141)$$

In contrast to (138), there does not seem to be a simple combinatorial derivation of (141). (Note that the first equality of (138) together with either equality of (141) gives the other equality of (141).)

It can be seen that (141), together with (104), provides an identity (referred to after (90)) involving $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ and $Z_n^{\mathrm{quad}}(x,y;z_3,z_2,z_1,z_4)$ (or $Z_n^{\mathrm{quad}}(x,y;z_1,z_2,z_3,z_4)$ and $Z_n^{\mathrm{quad}}(x,y;z_1,z_4,z_3,z_2)$).

It follows from (133) and (136), together with (130)–(131), (135) and (137)–(138), that

$$\begin{multline}
y^7(z_4-z_2)\,(z_1-z_3)\,Y_n(x,y;z_1,z_2,z_3,z_4)\,Z_{n-2}(x,y)=\\
\bigl(x(z_1-1)(z_2-1)-y\bigr)\bigl((z_1-1)(z_2-1)-yz_1z_2\bigr)\bigl(x(z_3-1)(z_4-1)-y\bigr)\times\\
\bigl((z_3-1)(z_4-1)-yz_3z_4\bigr)\,Y_{n-1}(x,y;z_1,z_4)\,Y_{n-1}(x,y;z_2,z_3)\,-\\
\bigl(x(z_4-1)(z_1-1)-y\bigr)\bigl((z_4-1)(z_1-1)-yz_4z_1\bigr)\bigl(x(z_2-1)(z_3-1)-y\bigr)\times\\
\bigl((z_2-1)(z_3-1)-yz_2z_3\bigr)\,Y_{n-1}(x,y;z_1,z_2)\,Y_{n-1}(x,y;z_3,z_4).\quad(142)
\end{multline}$$

This can be regarded as a restatement of (79), where (104) and (136) enable the conversion between (79) and (142).

Setting $z_2=1$ in (142), and relabeling z_3 as z_2 and z_4 as z_3, gives

$$\begin{multline}
y(z_1-z_2)\,(z_3-1)\,Y_n(x,y;z_1,1,z_2,z_3)\,Z_{n-2}(x,y)=\\
\bigl(x(z_2-1)(z_3-1)-y\bigr)\bigl((z_2-1)(z_3-1)-yz_2z_3\bigr)z_1\,Y_{n-1}(x,y;z_1,z_3)\,Z_{n-1}(x,y;z_2)\,-\\
\bigl(x(z_3-1)(z_1-1)-y\bigr)\bigl((z_3-1)(z_1-1)-yz_3z_1\bigr)z_2\,Y_{n-1}(x,y;z_2,z_3)\,Z_{n-1}(x,y;z_1),\quad(143)
\end{multline}$$

which can be regarded as a restatement of (81).

Setting $z_3=0$ in (142), using (139), replacing n by $n+1$, and relabeling z_3 as z_2 and z_4 as z_3 gives

$$\begin{aligned}&y^2(z_1-z_2)\,z_3\,Y_n(x,y;z_1,1,z_2,z_3)\,Z_{n-1}(x,y)=\\&\quad(z_1-1)\big(x(z_2-1)(z_3-1)-y\big)\big((z_2-1)(z_3-1)-yz_2z_3\big)\,Y_n(x,y;z_1,z_3)\,Z_{n-1}(x,y;z_2)-\\&\qquad(z_2-1)\big(x(z_1-1)(z_3-1)-y\big)\big((z_1-1)(z_3-1)-yz_1z_3\big)\,Y_n(x,y;z_2,z_3)\,Z_{n-1}(x,y;z_1),\end{aligned}\tag{144}$$

which can be regarded as a restatement of (84).

It can be seen that various other identities from Section 4 involving quadruply-refined, triply-refined or adjacent-boundary doubly-refined ASM generating functions could similarly be restated in terms of the functions (134)–(135).

Finally, note that some further quadratic identities satisfied by ASM generating functions can be obtained from (142), or by combining certain variations of the Desnanot–Jacobi identity with the Izergin–Korepin formula, for certain assignments of the spectral parameters, and then using (140).

An example of such an identity is

$$\begin{aligned}&(z_1-z_2)\,(z_3-z_4)\,Y_n(x,y;z_1,w_1,z_2,w_2)\,Y_n(x,y;z_3,w_1,z_4,w_2)-\\&\qquad(z_1-z_3)\,(z_2-z_4)\,Y_n(x,y;z_1,w_1,z_3,w_2)\,Y_n(x,y;z_2,w_1,z_4,w_2)+\\&\qquad\quad(z_1-z_4)\,(z_2-z_3)\,Y_n(x,y;z_1,w_1,z_4,w_2)\,Y_n(x,y;z_2,w_1,z_3,w_2)=0.\end{aligned}\tag{145}$$

This reduces to (60) for $w_1=w_2=1$, and can be obtained by applying (142) to each of the six cases of $(z_i-z_j)Y_n(x,y;z_i,w_1,z_j,w_2)$ on the LHS, and then checking that the overall expression on the LHS vanishes.

5.8. **Derivations of (19) and (20).**

In this section, the explicit expressions (19) and (20) for the ASM generating functions at $y=x+1$ are derived using a method different from that used in Section 3.2, which instead involves the six-vertex model with DWBC.

Let the crossing parameter q, for the weights (121), be given by $q=\pm i$, this corresponding to the so-called free fermion case of the six-vertex model. It can be shown that, for this assignment of q, the partition function (113), with weights given by (114) and (121), is explicitly

$$\begin{aligned}&Z_n^{6\mathrm{V}}(u_1,\ldots,u_n;u_{n+1},\ldots,u_{2n})\big|_{q=\pm i}=\\&\qquad(\pm2)^n\,i^{n^2}\,\textstyle\prod_{i=1}^n u_i^{1/2}v_i^{1/2}\,\prod_{1\le i<j\le n}(u_i+u_j)(v_i+v_j).\end{aligned}\tag{146}$$

This result can be obtained by combining the Izergin–Korepin formula (123) (in which $q=\pm i$ gives $a(u,v)b(u,v)=-(u^2+v^2)$) with the Cauchy double alternant evaluation

$$\det_{1\le i,j\le n}\left(\frac{1}{\alpha_i+\beta_j}\right)=\frac{\prod_{1\le i<j\le n}(\alpha_i-\alpha_j)(\beta_i-\beta_j)}{\prod_{1\le i,j\le n}(\alpha_i+\beta_j)}.\tag{147}$$

For previous appearances of (146), see, for example, Bogoliubov, Pronko and Zvonarev [20, Eq. (58)], or Okada [111, Thm. 2.4(1), third eq.], and for information regarding (147), see, for example, Muir [108, p. 345], [109, Sec. 353].

Assigning the spectral parameters in (146) according to (115), applying the relation (120), and using the first ten paramaterizations of (126), which enable arbitrary x and $z_1,\dots,z_4$ to be obtained, with y, z_{41}, z_{12}, z_{23} and z_{34} then being given by $y=x+1$ and (130), it follows that

$$\begin{aligned}
&Z_n^{\mathrm{gen}}\bigl(x,x+1;z_1,z_2,z_3,z_4;1-\tfrac{(z_4-1)(z_1-1)}{(x+1)\,z_4\,z_1},1-\tfrac{x(z_1-1)(z_2-1)}{x+1},1-\tfrac{(z_2-1)(z_3-1)}{(x+1)\,z_2\,z_3},1-\tfrac{x(z_3-1)(z_4-1)}{x+1}\bigr)\\
&\qquad\qquad=(xz_1z_3+1)(xz_2z_4+1)\bigl((xz_1+1)(xz_2+1)(xz_3+1)(xz_4+1)\bigr)^{n-2}\times\\
&\qquad\qquad\qquad\qquad\qquad\qquad(x+1)^{(n-2)(n-3)/2-2n+3}. \qquad (148)
\end{aligned}$$

(Using (136), the function in (148) could be written as $(x+1)^{-4}Y_n(x,x+1;z_1,z_2,z_3,z_4)$.)

The required expressions (19)–(20) now follow from (148) by applying (104)–(105), and, in some cases, setting boundary parameters to 1.

5.9. **Derivation of** (71).

In this section, a derivation is given of the identity (71) satisfied by the ASM generating function (66) associated with several rows (or several columns), for certain assignments of some of its parameters.

For $0\le m\le n$ and $1\le k_1<\ldots<k_m\le n$, and indeterminates r, s and $t_1,\dots,t_m$, consider the partition function $Z_n^{\mathrm{6V}}(r,\dots,r,t_1,r,\dots\dots,r,t_m,r,\dots,r;s,\dots\dots,s)$, as given by (113)–(114), with arbitrary functions a, b and c, where t_i appears in position k_i within $r,\dots,r,t_1,r,\dots\dots,r,t_m,r,\dots,r$. Using the bijection of Section 5.2 between $\mathrm{ASM}(n)$ and $\mathrm{6VDW}(n)$, and the behavior (111) of the relevant statistics under this bijection, it can be checked (by applying a process similar to that used in Section 5.2 for the derivation of (120)) that this partition function is related to the ASM generating function of (66) by

$$\begin{aligned}
&Z_n^{\mathrm{6V}}(r,\dots,r,t_1,r,\dots\dots,r,t_m,r,\dots,r;s,\dots\dots,s)=\\
&\quad b(r,s)^{(n-1)(n-m)}\,c(r,s)^{n-m}\,\textstyle\prod_{i=1}^m b(t_i,s)^{n-1}\,c(t_i,s)\times\\
&\qquad Z_n^{k_1,\dots,k_m}\Bigl(\bigl(\tfrac{a(r,s)}{b(r,s)}\bigr)^2,\bigl(\tfrac{c(r,s)}{b(r,s)}\bigr)^2;\ \tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\dots,\tfrac{a(t_m,s)\,b(r,s)}{a(r,s)\,b(t_m,s)};\ \bigl(\tfrac{c(t_1,s)}{b(t_1,s)}\bigr)^2,\dots,\bigl(\tfrac{c(t_m,s)}{b(t_m,s)}\bigr)^2\Bigr). \qquad (149)
\end{aligned}$$

(Note that the case of (120) obtained by setting $t_2=t_4=s$, and then relabeling t_3 as t_2, matches the case $m=2$, $k_1=1$ and $k_2=n$ of (149).)

It follows from a result of Colomo and Pronko [47, Eq. (6.8)], [48, Eq. (A.13)] (with a special case stated previously by Colomo and Pronko [46, Eq. 5.8]) that if the functions a, b and c are given by (121) (which involves the further indeterminate q), then, for $1\le m\le n$,

$$\begin{aligned}
&\bigl(a(r,s)\,b(r,s)\bigr)^{m(m-1)/2}\,\textstyle\prod_{1\le i<j\le m}(t_i-t_j)\times\\
&\quad Z_n^{\mathrm{6V}}(r,\dots,r,t_1,r,\dots\dots,r,t_m,r,\dots,r;s,\dots\dots,s)\,\textstyle\prod_{i=1}^{m-1}Z_{n-i}^{\mathrm{6V}}(r,\dots,r;s,\dots,s)\\
&\qquad\qquad=\det_{1\le i,j\le m}\Bigl(\bigl(a(t_i,s)\,b(t_i,s)\bigr)^{j-1}\,(t_i-r)^{m-j}\,Z_{n-j+1}^{\mathrm{6V}}(t_i,r,\dots,r;s,\dots\dots,s)\Bigr), \qquad (150)
\end{aligned}$$

where the positions of $t_1,\dots,t_m$ within $r,\dots,r,t_1,r,\dots\dots,r,t_m,r,\dots,r$ on the LHS of (150) are now immaterial. An alternative proof of (150), in which a general identity for minors of a matrix is combined with the Izergin–Korepin formula (123), will be given elsewhere by this author. A result related to (150), in the case $q=e^{\pm 2\pi i/3}$, has been obtained by Ayyer and Romik [12, Thm. 5].

Using (149) to express each case of a partition function in (150) as a case of the ASM generating function (66), and applying the definition (69), gives

$$Z_n^{k_1,\dots,k_m}\Bigl(\bigl(\tfrac{a(r,s)}{b(r,s)}\bigr)^2,\bigl(\tfrac{c(r,s)}{b(r,s)}\bigr)^2;\ \tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\dots,\tfrac{a(t_m,s)\,b(r,s)}{a(r,s)\,b(t_m,s)};\ \bigl(\tfrac{c(t_1,s)}{b(t_1,s)}\bigr)^2,\dots,\bigl(\tfrac{c(t_m,s)}{b(t_m,s)}\bigr)^2\Bigr)=\\ X_n\Bigl(\bigl(\tfrac{a(r,s)}{b(r,s)}\bigr)^2,\bigl(\tfrac{c(r,s)}{b(r,s)}\bigr)^2;\ \tfrac{a(t_1,s)\,b(r,s)}{a(r,s)\,b(t_1,s)},\dots,\tfrac{a(t_m,s)\,b(r,s)}{a(r,s)\,b(t_m,s)}\Bigr).\qquad(151)$$

The required result (71) follows immediately from (151) by parameterizing x, y, $z_1,\dots,z_m$ and $w_1,\dots,w_m$ in terms of q, r, s and $t_1,\dots,t_m$ as $x=\bigl(\frac{a(r,s)}{b(r,s)}\bigr)^2$, $y=\bigl(\frac{c(r,s)}{b(r,s)}\bigr)^2$, $z_i=\frac{a(t_i,s)\,b(r,s)}{a(r,s)\,b(t_i,s)}$ and $w_i=\bigl(\frac{c(t_i,s)}{b(t_i,s)}\bigr)^2$, and observing that this parameterization enables arbitrary x, y and $z_1,\dots,z_m$ to be obtained, with $w_1,\dots,w_m$ then being given by $w_i=xz_i^2+(y-x-1)z_i+1$.

5.10. **Derivations of (48) and (73).** In this section, an expression for the partition function of the six-vertex model with DWBC at a certain value of its crossing parameter is given. This result, together with certain other previously-stated results, is then used to obtain derivations of the identity (48) satisfied by the alternative quadruply-refined ASM generating function at $x=y=1$, and of the identity (73) satisfied by the function (69) at $x=y=1$.

Let the crossing parameter q, for the weights (121), be given by $q=e^{\pm 2\pi i/3}$, this corresponding to the so-called combinatorial point of the six-vertex model. It can be shown that, for this assignment of q, the partition function (113), with weights given by (114) and (121), can be expressed as

$$Z_n^{\mathrm{6V}}(u_1,\dots,u_n;u_{n+1},\dots,u_{2n})\big|_{q=e^{\pm 2\pi i/3}}=\\ (\pm 1)^n\, i^{n^2}\, 3^{n/2}\, u_1^{1/2}\dots u_{2n}^{1/2}\ s_{(n-1,n-1,\dots,2,2,1,1)}(u_1,\dots,u_{2n}),\qquad(152)$$

where $s_{(n-1,n-1,\dots,2,2,1,1)}(u_1,\dots,u_{2n})$ is the Schur function indexed by the double-staircase partition $(n-1,n-1,\dots,2,2,1,1)$, evaluated at the spectral parameters $u_1,\dots,u_{2n}$.

An important consequence of (152) is that $Z_n^{\mathrm{6V}}(u_1,\dots,u_n;u_{n+1},\dots,u_{2n})\big|_{q=e^{\pm 2\pi i/3}}$ is symmetric in all spectral parameters $u_1,\dots,u_{2n}$.

The result (152) was first obtained by Okada [111, Thm. 2.4(1), second equation], and Stroganov [140, Eq. (17)]. For further related information, derivations and results, see, for example, Aval [7], Fonseca and Zinn-Justin [75, Sec. B.2], Lascoux [98, p. 4], Razumov and Stroganov [117, Sec. 2], or Zinn-Justin [149, Sec. 2.5.6].

Proceeding to the derivation of (48), using the symmetry of the LHS of (152) in all of its spectral parameters, it follows that

$$Z_n^{6\mathrm{V}}(t_1,1,\ldots,1,t_3;t_4,1,\ldots,1,t_2)\big|_{q=e^{\pm 2\pi i/3}} = \\ Z_n^{6\mathrm{V}}(t_1,t_2,t_3,t_4,1,\ldots,1;1,\ldots\ldots,1)\big|_{q=e^{\pm 2\pi i/3}}. \qquad (153)$$

Applying (120) and (149) (with $r=s=1$, $m=4$ and $k_i=i$) to the LHS and RHS, respectively, of (153), and then using (121) (with $q=e^{\pm 2\pi i/3}$) for the functions a, b and c (and noting that $a(1,1)=b(1,1)$ and $b(1,1)^2=c(1,1)^2$), gives

$$\left(\tfrac{b(1,t_2)\,b(1,t_4)}{b(t_2,1)\,b(t_4,1)}\right)^{n-1} Z_n^{\mathrm{gen}}\Bigl(1,1;\ \tfrac{a(t_1,1)}{b(t_1,1)},\tfrac{a(1,t_2)}{b(1,t_2)},\tfrac{a(t_3,1)}{b(t_3,1)},\tfrac{a(1,t_4)}{b(1,t_4)}; \\ \tfrac{a(1,1)\,a(t_1,t_4)}{a(t_1,1)\,a(1,t_4)},\tfrac{b(1,1)\,b(t_1,t_2)}{b(t_1,1)\,b(1,t_2)},\tfrac{a(1,1)\,a(t_3,t_2)}{a(t_3,1)\,a(1,t_2)},\tfrac{b(1,1)\,b(t_3,t_4)}{b(t_3,1)\,b(1,t_4)}\Bigr)\Big|_{q=e^{\pm 2\pi i/3}} =$$

$$Z_n^{1,2,3,4}\Bigl(1,1;\ \tfrac{a(t_1,1)}{b(t_1,1)},\tfrac{a(t_2,1)}{b(t_2,1)},\tfrac{a(t_3,1)}{b(t_3,1)},\tfrac{a(t_4,1)}{b(t_4,1)};\ \left(\tfrac{c(t_1,1)}{b(t_1,1)}\right)^2,\left(\tfrac{c(t_2,1)}{b(t_2,1)}\right)^2,\left(\tfrac{c(t_3,1)}{b(t_3,1)}\right)^2,\left(\tfrac{c(t_4,1)}{b(t_4,1)}\right)^2\Bigr)\Big|_{q=e^{\pm 2\pi i/3}}. \qquad (154)$$

It follows from (154), by applying (151) to the RHS, and then setting $t_i=\frac{qz_i+1}{z_i+q}$, or equivalently $z_i=\frac{qt_i-1}{q-t_i}=\frac{a(t_i,1)}{b(t_i,1)}$, for each $1\le i\le 4$, that

$$(z_2z_4)^{n-1}Z_n^{\mathrm{gen}}\Bigl(1,1;z_1,\tfrac{1}{z_2},z_3,\tfrac{1}{z_4};1+\tfrac{(z_4-1)(z_1-1)}{z_1},1+\tfrac{(z_1-1)(z_2-1)}{z_2},1+\tfrac{(z_2-1)(z_3-1)}{z_3},1+\tfrac{(z_3-1)(z_4-1)}{z_4}\Bigr) \\ = X_n(1,1;z_1,z_2,z_3,z_4). \qquad (155)$$

The required result (48) now follows from (155) by applying (104) to the LHS, and using the first definition of (10), the definition (69) and the identity (89).

Proceeding to the derivation of (73), setting $r=s=1$, $q=e^{\pm 2\pi i/3}$ and $k_i=i$ in (150), and then applying (152) to the partition function $Z_n^{6\mathrm{V}}(t_1,\ldots,t_m,1,\ldots,1;1,\ldots,1)$, applying (149) (with $m=0$ or $m=1$) to the remaining cases of partition functions, and using (69) and (121), gives

$$s_{(n-1,n-1,\ldots,2,2,1,1)}(t_1,\ldots,t_m,\underbrace{1,\ldots,1}_{2n-m}) = \\ 3^{n(n-1)/2}\,b(1,1)^{-m(n-1)}\,(b(t_1,1)\ldots b(t_m,1))^{n-1}\,X_n\bigl(1,1;\ \tfrac{a(t_1,1)}{b(t_1,1)},\ldots,\tfrac{a(t_m,1)}{b(t_m,1)}\bigr)\big|_{q=e^{\pm 2\pi i/3}}. \qquad (156)$$

(Alternatively, this can be obtained by combining (121), (149), (151) and (152).)

The required result (73) now follows from (156) by setting $t_i=\frac{qz_i+1}{z_i+q}$, or equivalently $z_i=\frac{qt_i-1}{q-t_i}=\frac{a(t_i,1)}{b(t_i,1)}$, for each $1\le i\le m$.

REFERENCES

[1] F. Abeles, *Dodgson condensation: the historical and mathematical development of an experimental method*, Linear Algebra Appl. **429** (2008), no. 2–3, 429–438, doi. MR2419937

[2] G. Andrews, *Plane partitions. III. The weak Macdonald conjecture*, Invent. Math. **53** (1979), no. 3, 193–225, doi. MR549398

[3] ______, *Macdonald's conjecture and descending plane partitions*, Combinatorics, representation theory and statistical methods in groups, Lecture Notes in Pure and Appl. Math., vol. 57, Dekker, New York, 1980, pp. 91–106. MR588196

[4] ______, *Plane partitions. V. The TSSCPP conjecture*, J. Combin. Theory Ser. A **66** (1994), no. 1, 28–39, `doi`. MR1273289

[5] G. Andrews and D. Stanton, *Determinants in plane partition enumeration*, European J. Combin. **19** (1998), no. 3, 273–282, `doi`. MR1621001

[6] J.-C. Aval, *Keys and alternating sign matrices*, Sém. Lothar. Combin. **59** (2007/10), Art. B59f, 13pp. (electronic), `arXiv:0711.2150`. MR2465402

[7] ______, *On the symmetry of the partition function of some square ice models*, Theoret. and Math. Phys. **161** (2009), no. 3, 1582–1589, `arXiv:0903.0777`, `doi`. MR2642191

[8] J.-C. Aval and P. Duchon, *Enumeration of alternating sign matrices of even size (quasi)-invariant under a quarter-turn rotation*, 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), Discrete Math. Theor. Comput. Sci. Proc., AK, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2009, pp. 115–126, `arXiv:0906.3445`. MR2721506

[9] ______, *Enumeration of alternating sign matrices of even size (quasi-)invariant under a quarter-turn rotation*, Electron. J. Combin. **17** (2010), no. 1, Research Paper 51, 20pp. (electronic), `arXiv:0910.3047`. MR2607337

[10] A. Ayyer, *A natural bijection between permutations and a family of descending plane partitions*, European J. Combin. **31** (2010), no. 7, 1785–1791, `arXiv:0909.4732`, `doi`. MR2673018

[11] A. Ayyer, R. Cori, and D. Gouyou-Beauchamps, *Monotone triangles and 312 pattern avoidance*, Electron. J. Combin. **18** (2011), no. 2, Paper 26, 22 pp. (electronic), `arXiv:1101.1666`. MR2861418

[12] A. Ayyer and D. Romik, *New enumeration formulas for alternating sign matrices and square ice partition functions*, 2012, preprint, `arXiv:1202.3651`.

[13] M. T. Batchelor, J. de Gier, and B. Nienhuis, *The quantum symmetric XXZ chain at $\Delta = -\frac{1}{2}$, alternating-sign matrices and plane partitions*, J. Phys. A **34** (2001), no. 19, L265–L270, `arXiv:cond-mat/0101385`, `doi`. MR1836155

[14] R. Baxter, *Exactly solved models in statistical mechanics*, Academic Press Inc. [Harcourt Brace Jovanovich Publishers], London, 1989, Reprint of the 1982 original. MR998375

[15] R. Behrend, *Osculating paths and oscillating tableaux*, Electron. J. Combin. **15** (2008), no. 1, Research Paper 7, 56 pp. (electronic), `arXiv:math/0701755`. MR2368912

[16] R. Behrend, P. Di Francesco, and P. Zinn-Justin, *A doubly-refined enumeration of alternating sign matrices and descending plane partitions*, 2012, To appear in J. Combin. Theory Ser. A, `arXiv:1202.1520`.

[17] ______, *On the weighted enumeration of alternating sign matrices and descending plane partitions*, J. Combin. Theory Ser. A **119** (2012), no. 2, 331–363, `arXiv:1103.1176`, `doi`. MR2860598

[18] P. Biane, L. Cantini, and A. Sportiello, *Doubly-refined enumeration of alternating sign matrices and determinants of 2-staircase Schur functions*, 2011, preprint, `arXiv:1101.3427`.

[19] N. Bogoliubov, A. Kitaev, and M. Zvonarev, *Boundary polarization in the six-vertex model*, Phys. Rev. E (3) **65** (2002), no. 2, 026126, 4, `arXiv:cond-mat/0107146`, `doi`. MR1922214

[20] N. Bogoliubov, A. Pronko, and M. Zvonarev, *Boundary correlation functions of the six-vertex model*, J. Phys. A **35** (2002), no. 27, 5525–5541, `arXiv:math-ph/0203025`, `doi`. MR1917248

[21] M. Bousquet-Mélou and L. Habsieger, *Sur les matrices à signes alternants*, Discrete Math. **139** (1995), no. 1-3, 57–72, Formal power series and algebraic combinatorics (Montreal, PQ, 1992), `doi`. MR1336832

[22] D. Bressoud, *Proofs and confirmations. the story of the alternating sign matrix conjecture*, MAA Spectrum, Mathematical Association of America and Cambridge University Press, Washington, DC and Cambridge, 1999. MR1718370

[23] ———, *Three alternating sign matrix identities in search of bijective proofs*, Adv. in Appl. Math. **27** (2001), no. 2-3, 289–297, Special issue in honor of Dominique Foata's 65th birthday (Philadelphia, PA, 2000), `arXiv:math/0007114`, `doi`. MR1868966

[24] ———, *Exploiting symmetries: alternating-sign matrices and the Weyl character formulas*, Surveys in number theory, Dev. Math., vol. 17, Springer, New York, 2008, pp. 43–55. MR2462945

[25] D. Bressoud and J. Propp, *How the alternating sign matrix conjecture was solved*, Notices Amer. Math. Soc. **46** (1999), no. 6, 637–646. MR1691562

[26] R. Brualdi and S. Kirkland, *Aztec diamonds and digraphs, and Hankel determinants of Schröder numbers*, J. Combin. Theory Ser. B **94** (2005), no. 2, 334–351, `doi`. MR2145518

[27] R. Brualdi and H. Schneider, *Determinantal identities: Gauss, Schur, Cauchy, Sylvester, Kronecker, Jacobi, Binet, Laplace, Muir, and Cayley*, Linear Algebra Appl. **52/53** (1983), 769–791, `doi`. MR1500275

[28] B. Brubaker, D. Bump, and S. Friedberg, *Eisenstein series, crystals, and ice*, Notices Amer. Math. Soc. **58** (2011), no. 11, 1563–1571. MR2896085

[29] ———, *Schur polynomials and the Yang-Baxter equation*, Comm. Math. Phys. **308** (2011), no. 2, 281–301, `arXiv:0912.0911`, `doi`. MR2851143

[30] D. Bump, P. McNamara, and M. Nakasuji, *Factorial Schur functions and the Yang-Baxter equation*, 2011, preprint, `arXiv:1108.3087`.

[31] L. Cantini and A. Sportiello, *Proof of the Razumov-Stroganov conjecture*, J. Combin. Theory Ser. A **118** (2011), no. 5, 1549–1574, `arXiv:1003.3376`, `doi`. MR2771600

[32] ———, *A one-parameter refinement of the Razumov-Stroganov correspondence*, 2012, preprint, `arXiv:1202.5253`.

[33] R. Chapman, *Alternating sign matrices and tournaments*, Adv. in Appl. Math. **27** (2001), no. 2-3, 318–335, Special issue in honor of Dominique Foata's 65th birthday (Philadelphia, PA, 2000), `arXiv:math/0008029`, `doi`. MR1868968

[34] H. Cheballah, *Combinatoire des matrices à signes alternants et des partitions planes*, 2011, PhD thesis, Université Paris Nord, `http://www-lipn.univ-paris13.fr/~cheballah/memoires/these.pdf`.

[35] H. Cheballah and P. Biane, *Gog and magog triangles, and the Schützenberger involution*, Sém. Lothar. Combin. **66** (2012), Art. B66d, 20pp. (electronic), `arXiv:1105.4986`.

[36] M. Ciucu, *Perfect matchings of cellular graphs*, J. Algebraic Combin. **5** (1996), no. 2, 87–103, `doi`. MR1382040

[37] ———, *A complementation theorem for perfect matchings of graphs having a cellular completion*, J. Combin. Theory Ser. A **81** (1998), no. 1, 34–68, `doi`. MR1492868

[38] M. Ciucu, T. Eisenkölbl, C. Krattenthaler, and D. Zare, *Enumeration of lozenge tilings of hexagons with a central triangular hole*, J. Combin. Theory Ser. A **95** (2001), no. 2, 251–334, `arXiv:math/9912053`, `doi`. MR1845144

[39] M. Ciucu and C. Krattenthaler, *Plane partitions. II. $5\frac{1}{2}$ symmetry classes*, Combinatorial methods in representation theory (Kyoto, 1998), Adv. Stud. Pure Math., vol. 28, Kinokuniya, Tokyo, 2000, pp. 81–101, `arXiv:math/9808018`. MR1855591

[40] F. Colomo, Personal correspondence, March 2012.

[41] F. Colomo and A. Pronko, *On some representations of the six vertex model partition function*, Phys. Lett. A **315** (2003), no. 3–4, 231–236, `doi`. MR2003060

[42] ———, *On the partition function of the six-vertex model with domain wall boundary conditions*, J. Phys. A **37** (2004), no. 6, 1987–2002, `arXiv:math-ph/0309064`, `doi`. MR2045907

[43] ———, *On the refined 3-enumeration of alternating sign matrices*, Adv. in Appl. Math. **34** (2005), no. 4, 798–811, `arXiv:math-ph/0404045`, `doi`. MR2128998

[44] ______, *On two-point boundary correlations in the six-vertex model with domain wall boundary conditions*, J. Stat. Mech. Theory Exp. (2005), P05010, 21 pp. (electronic), `arXiv:math-ph/0503049`, doi. MR2149859

[45] ______, *Square ice, alternating sign matrices, and classical orthogonal polynomials*, J. Stat. Mech. Theory Exp. (2005), P01005, 33 pp. (electronic), `arXiv:math-ph/0411076`, doi. MR2114554

[46] ______, *The role of orthogonal polynomials in the six-vertex model and its combinatorial applications*, J. Phys. A **39** (2006), no. 28, 9015–9033, `arXiv:math-ph/0602033`, doi. MR2240471

[47] ______, *Emptiness formation probability in the domain-wall six-vertex model*, Nuclear Phys. B **798** (2008), no. 3, 340–362, `arXiv:0712.1524`, doi. MR2411855

[48] ______, *The arctic curve of the domain-wall six-vertex model*, J. Stat. Phys. **138** (2010), no. 4-5, 662–700, `arXiv:0907.1264`, doi. MR2594917

[49] ______, *An approach for calculating correlation functions in the six-vertex model with domain wall boundary conditions*, Theoret. and Math. Phys. **171** (2012), no. 2, 641–654, `arXiv:1111.4353`, doi.

[50] R. Cori, P. Duchon, and F. Le Gac, *Comptage de matrices à signe alternant en fonction du nombre d'entrées négatives*, 2010, `http://www-apr.lip6.fr/alea2010/transparents/le_gac.pdf`.

[51] J. de Gier, *Loops, matchings and alternating-sign matrices*, Discrete Math. **298** (2005), no. 1-3, 365–388, `arXiv:math/0211285`, doi. MR2163456

[52] ______, *The Razumov-Stroganov conjecture: stochastic processes, loops and combinatorics*, J. Stat. Mech. Theory Exp. (2007), N02001, 6 pp. (electronic), `arXiv:cond-mat/0702432`, doi. MR2293822

[53] ______, *Fully packed loop models on finite geometries*, Polygons, polyominoes and polycubes, Lecture Notes in Phys., vol. 775, Springer, Dordrecht, 2009, pp. 317–346, `arXiv:0901.3963`. MR2790336

[54] J. de Gier, P. Pyatov, and P. Zinn-Justin, *Punctured plane partitions and the q-deformed Knizhnik–Zamolodchikov and Hirota equations*, J. Combin. Theory Ser. A **116** (2009), no. 4, 772–794, `arXiv:0712.3584`, doi. MR2513634

[55] P. Di Francesco, *A refined Razumov–Stroganov conjecture. II*, J. Stat. Mech. Theory Exp. (2004), P11004, 20 pp. (electronic), `arXiv:cond-mat/0409576`, doi. MR2110544

[56] ______, *A refined Razumov-Stroganov conjecture*, J. Stat. Mech. Theory Exp. (2004), P08009, 16 pp. (electronic), `arXiv:cond-mat/0407477`, doi. MR2115252

[57] ______, *Totally symmetric self-complementary plane partitions and the quantum Knizhnik-Zamolodchikov equation: a conjecture*, J. Stat. Mech. Theory Exp. (2006), P09008, 14 pp. (electronic), `arXiv:cond-mat/0607499`, doi. MR2278472

[58] ______, *Open boundary quantum Knizhnik-Zamolodchikov equation and the weighted enumeration of plane partitions with symmetries*, J. Stat. Mech. Theory Exp. (2007), P01024, 22 pp. (electronic), `arXiv:math-ph/0611012`, doi. MR2284008

[59] P. Di Francesco and P. Zinn-Justin, *Around the Razumov–Stroganov conjecture: proof of a multi-parameter sum rule*, Electron. J. Combin. **12** (2005), Research Paper 6, 27 pp. (electronic), `arXiv:math-ph/0410061`. MR2134169

[60] C. Dodgson, *Condensation of determinants, being a new and brief method for computing their arithmetical values*, Proc. Roy. Soc. London **15** (1866), 150–155, doi.

[61] W. Doran, IV, *A connection between alternating sign matrices and totally symmetric self-complementary plane partitions*, J. Combin. Theory Ser. A **64** (1993), no. 2, doi. MR1245163

[62] Ö. Eğecioğlu, T. Redmond, and C. Ryavec, *From a polynomial Riemann hypothesis to alternating sign matrices*, Electron. J. Combin. **8** (2001), no. 1, Research Paper 36, 51 pp. (electronic), `http://www.combinatorics.org/Volume_8/Abstracts/v8i1r36.html`. MR1877655

[63] N. Elkies, G. Kuperberg, M. Larsen, and J. Propp, *Alternating-sign matrices and domino tilings. I*, J. Algebraic Combin. **1** (1992), no. 2, 111–132, `arXiv:math/9201305`, doi. MR1226347

[64] ______, *Alternating-sign matrices and domino tilings. II*, J. Algebraic Combin. **1** (1992), no. 3, 219–234, `arXiv:math/9201305`, doi. MR1194076

[65] S. Eu and T. Fu, *A simple proof of the Aztec diamond theorem*, Electron. J. Combin. **12** (2005), Research Paper 18, 8 pp. (electronic), `arXiv:math/0412041`. MR2134181

[66] P. Ferrari and H. Spohn, *Domino tilings and the six-vertex model at its free-fermion point*, J. Phys. A **39** (2006), no. 33, 10297–10306, `arXiv:cond-mat/0605406`, `doi`. MR2256593

[67] I. Fischer, *The number of monotone triangles with prescribed bottom row*, Adv. in Appl. Math. **37** (2006), no. 2, 249–267, `arXiv:math/0501102`, `doi`. MR2251438

[68] ———, *A new proof of the refined alternating sign matrix theorem*, J. Combin. Theory Ser. A **114** (2007), no. 2, 253–264, `arXiv:math/0507270`, `doi`. MR2293090

[69] ———, *The operator formula for monotone triangles—simplified proof and three generalizations*, J. Combin. Theory Ser. A **117** (2010), no. 8, 1143–1157, `arXiv:0903.4628`, `doi`. MR2677680

[70] ———, *Refined enumerations of alternating sign matrices: monotone (d, m)-trapezoids with prescribed top and bottom row*, J. Algebraic Combin. **33** (2011), no. 2, 239–257, `arXiv:0907.0401`, `doi`. MR2765324

[71] ———, *Linear relations of refined enumerations of alternating sign matrices*, J. Combin. Theory Ser. A **119** (2012), no. 3, 556–578, `arXiv:1008.0527`, `doi`.

[72] I. Fischer and D. Romik, *More refined enumerations of alternating sign matrices*, Adv. Math. **222** (2009), no. 6, 2004–2035, `arXiv:0903.5073`, `doi`. MR2562772

[73] O. Foda and I. Preston, *On the correlation functions of the domain wall six-vertex model*, J. Stat. Mech. Theory Exp. (2004), P11001, 24 pp. (electronic), `arXiv:math-ph/0409067`, `doi`. MR2110546

[74] T. Fonseca, *Matrices à signes alternants, boucles denses et partitions planes*, 2010, PhD thesis, Université Pierre et Marie Curie, Paris VI, `http://tel.archives-ouvertes.fr/docs/00/53/72/16/PDF/tese.pdf`.

[75] T. Fonseca and P. Zinn-Justin, *On the doubly refined enumeration of alternating sign matrices and totally symmetric self-complementary plane partitions*, Electron. J. Combin. **15** (2008), Research Paper 81, 35 pp. (electronic), `arXiv:0803.1595`. MR2411458

[76] I. Gessel and G. Xin, *The generating function of ternary trees and continued fractions*, Electron. J. Combin. **13** (2006), no. 1, Research Paper 53, 48 pp. (electronic), `arXiv:math/0505217`. MR2240759

[77] A. Hamel and R. King, *Bijective proofs of shifted tableau and alternating sign matrix identities*, J. Algebraic Combin. **25** (2007), no. 4, 417–458, `arXiv:math/0507479`, `doi`. MR2320371

[78] A. Hone, *Dodgson condensation, alternating signs and square ice*, Philos. Trans. R. Soc. Lond. Ser. A Math. Phys. Eng. Sci. **364** (2006), no. 1849, 3183–3198, `doi`. MR2317901

[79] M. Ishikawa, *On refined enumerations of totally symmetric self-complementary plane partitions I*, 2006, preprint, `arXiv:math/0602068`.

[80] ———, *On refined enumerations of totally symmetric self-complementary plane partitions II*, 2006, preprint, `arXiv:math/0606082`.

[81] A. Izergin, *Partition function of a six-vertex model in a finite volume*, Dokl. Akad. Nauk SSSR **297** (1987), no. 2, 331–333. MR919260

[82] A. Izergin, D. Coker, and V. Korepin, *Determinant formula for the six-vertex model*, J. Phys. A **25** (1992), no. 16, 4315–4334, `doi`. MR1181591

[83] M. Karklinsky and D. Romik, *A formula for a doubly refined enumeration of alternating sign matrices*, Adv. in Appl. Math. **45** (2010), no. 1, 28–35, `arXiv:0906.3405`, `doi`. MR2628784

[84] D. Knuth, *Overlapping Pfaffians*, Electron. J. Combin. **3** (1996), no. 2, Research Paper 5, 13 pp. (electronic), `arXiv:math/9503234`. MR1392490

[85] V. Korepin, *Calculation of norms of Bethe wave functions*, Comm. Math. Phys. **86** (1982), no. 3, 391–418, `doi`. MR677006

[86] V. Korepin and P. Zinn-Justin, *Thermodynamic limit of the six-vertex model with domain wall boundary conditions*, J. Phys. A **33** (2000), no. 40, 7053–7066, `arXiv:cond-mat/0004250`, `doi`. MR1792450

[87] C. Krattenthaler, *A gog-magog conjecture*, 1996, `http://www.mat.univie.ac.at/~kratt/artikel/magog.html`.

[88] ———, *Advanced determinant calculus*, Sém. Lothar. Combin. **42** (1999), Art. B42q, 67 pp. (electronic), The Andrews Festschrift (Maratea, 1998), `arXiv:math/9902004`. MR1701596

[89] ———, *Advanced determinant calculus: a complement*, Linear Algebra Appl. **411** (2005), 68–166, `arXiv:math/0503507`, `doi`. MR2178686

[90] ———, *Descending plane partitions and rhombus tilings of a hexagon with a triangular hole*, European J. Combin. **27** (2006), no. 7, 1138–1146, `arXiv:math/0310188`, `doi`. MR2259946

[91] E. Kuo, *Applications of graphical condensation for enumerating matchings and tilings*, Theoret. Comput. Sci. **319** (2004), no. 1-3, 29–57, `arXiv:math/0304090`, `doi`. MR2074946

[92] G. Kuperberg, *Another proof of the alternating-sign matrix conjecture*, Internat. Math. Res. Notices (1996), no. 3, 139–150, `arXiv:math/9712207`. MR1383754

[93] ———, *Symmetry classes of alternating-sign matrices under one roof*, Ann. of Math. (2) **156** (2002), no. 3, 835–866, `arXiv:math/0008184`, `doi`. MR1954236

[94] P. Lalonde, *q-enumeration of alternating sign matrices with exactly one* -1, Discrete Math. **256** (2002), no. 3, 759–773, LaCIM 2000 Conference on Combinatorics, Computer Science and Applications (Montreal, QC), `doi`. MR1935787

[95] ———, *Lattice paths and the antiautomorphism of the poset of descending plane partitions*, Discrete Math. **271** (2003), no. 1–3, 311–319, `doi`. MR1999553

[96] ———, *Alternating sign matrices with one -1 under vertical reflection*, J. Combin. Theory Ser. A **113** (2006), no. 6, 980–994, `arXiv:math/0401339`, `doi`. MR2244128

[97] A. Lascoux, *The 6 vertex model and Schubert polynomials*, SIGMA Symmetry Integrability Geom. Methods Appl. **3** (2007), Paper 29, 12pp., `arXiv:math/0610719`, `doi`. MR2299830

[98] ———, *Gaudin functions, and Euler-Poincaré characteristics*, 2007, preprint, `arXiv:0709.1635`.

[99] F. Le Gac, *Comptage d'ASMs selon le nombre de* -1, 2011, `http://www.labri.fr/perso/marckert/LeGac.pdf`.

[100] B. Leclerc, *On identities satisfied by minors of a matrix*, Adv. Math. **100** (1993), no. 1, 101–132, `doi`. MR1224528

[101] P. McNamara, *Factorial Schur functions via the six vertex model*, 2009, preprint, `arXiv:0910.5288`.

[102] W. Mills, D. Robbins, and H. Rumsey, Jr., *Proof of the Macdonald conjecture*, Invent. Math. **66** (1982), no. 1, 73–87, `doi`. MR652647

[103] ———, *Alternating sign matrices and descending plane partitions*, J. Combin. Theory Ser. A **34** (1983), no. 3, 340–359, `doi`. MR700040

[104] ———, *Self-complementary totally symmetric plane partitions*, J. Combin. Theory Ser. A **42** (1986), no. 2, 277–292, `doi`. MR847558

[105] ———, *Enumeration of a symmetry class of plane partitions*, Discrete Math. **67** (1987), no. 1, 43–55, `doi`. MR908185

[106] K. Motegi, *Boundary correlation functions of the six and nineteen vertex models with domain wall boundary conditions*, Phys. A **390** (2011), no. 20, 3337–3347, `arXiv:1101.0187`, `doi`. MR2822902

[107] ———, *Boundary correlation functions of integrable vertex models*, J. Phys.: Conf. Ser. **343** (2012), no. 012081, 10 pp., `doi`.

[108] T. Muir, *The theory of determinants in the historical order of development. Vol. I*, Macmillan and Co., London, 1906.

[109] ———, *A treatise on the theory of determinants*, Revised and enlarged by William H. Metzler, Dover Publications Inc., New York, 1960. MR0114826

[110] S. Ng, *Alternating paths of fully packed loops and inversion number*, 2012, preprint, `arXiv:1202.2483`.

[111] S. Okada, *Enumeration of symmetry classes of alternating sign matrices and characters of classical groups*, J. Algebraic Combin. **23** (2006), no. 1, 43–69, `arXiv:math/0408234`, `doi`. MR2218849

[112] V. Pasquier, *Quantum incompressibility and Razumov–Stroganov type conjectures*, Ann. Henri Poincaré **7** (2006), no. 3, 397–421, `arXiv:cond-mat/0506075`, `doi`. MR2226742

[113] J. Propp, *The many faces of alternating-sign matrices*, Discrete models: combinatorics, computation, and geometry (Paris, 2001), Discrete Math. Theor. Comput. Sci. Proc., AA, Maison Inform. Math. Discrèt. (MIMD), Paris, 2001, pp. 43–58 (electronic), `arXiv:math/0208125`. MR1888762

[114] ———, *Lambda-determinants and domino-tilings*, Adv. in Appl. Math. **34** (2005), no. 4, 871–879, `arXiv:math/0406301`, `doi`. MR2129002

[115] A. Razumov and Y. Stroganov, *Spin chains and combinatorics*, J. Phys. A **34** (2001), no. 14, 3185–3190, `arXiv:cond-mat/0012141`, `doi`. MR1832783

[116] ———, *Combinatorial nature of the ground-state vector of the* O(1) *loop model*, Theoret. and Math. Phys. **138** (2004), no. 3, 333–337, `arXiv:math/0104216`, `doi`. MR2077318

[117] ———, *Refined enumerations of some symmetry classes of alternating-sign matrices*, Theoret. and Math. Phys. **141** (2004), no. 3, 1609–1630, `arXiv:math-ph/0312071`, `doi`. MR2141132

[118] ———, *Bethe roots and refined enumeration of alternating-sign matrices*, J. Stat. Mech. Theory Exp. (2006), P07004, 12 pp. (electronic), `arXiv:math-ph/0605004`, `doi`. MR2244324

[119] ———, *Enumeration of quarter-turn symmetric alternating-sign matrices of odd-order*, Theoret. and Math. Phys. **149** (2006), no. 3, 1639–1650, `arXiv:math-ph/0507003`, `doi`. MR2321099

[120] ———, *Enumerations of half-turn-symmetric alternating-sign matrices of odd-order*, Theoret. and Math. Phys. **148** (2006), no. 3, 1174–1198, `arXiv:math-ph/0504022`, `doi`. MR2283658

[121] ———, *A statistical model of three colors with boundary conditions of domain wall type: functional equations*, Theoret. and Math. Phys. **161** (2009), no. 1, 1325–1339, `arXiv:0805.0669`, `doi`. MR2664880

[122] A. Razumov, Y. Stroganov, and P. Zinn-Justin, *Polynomial solutions of qKZ equation and ground state of XXZ spin chain at* $\Delta = -1/2$, J. Phys. A **40** (2007), no. 39, 11827–11847, `arXiv:0704.3542`, `doi`. MR2374053

[123] D. Robbins, *The story of* $1, 2, 7, 42, 429, 7436, \ldots$, Math. Intelligencer **13** (1991), no. 2, 12–19, `doi`. MR1098216

[124] ———, *Symmetry classes of alternating sign matrices*, 2000, preprint, `arXiv:math/0008045`.

[125] D. Robbins and H. Rumsey, Jr., *Determinants and alternating sign matrices*, Adv. in Math. **62** (1986), no. 2, 169–184, `doi`. MR865837

[126] H. Rosengren, *An Izergin-Korepin-type identity for the 8VSOS model, with applications to alternating sign matrices*, Adv. in Appl. Math. **43** (2009), no. 2, 137–155, `arXiv:0801.1229`, `doi`. MR2531917

[127] ———, *The three-colour model with domain wall boundary conditions*, Adv. in Appl. Math. **46** (2011), no. 1-4, 481–535, `arXiv:0911.0561`, `doi`. MR2794035

[128] ———, *New proofs of determinant evaluations related to plane partitions*, 2012, preprint, `arXiv:1204.3424`.

[129] K. Sogo, *Time-dependent orthogonal polynomials and theory of soliton. Applications to matrix model, vertex model and level statistics*, J. Phys. Soc. Japan **62** (1993), no. 6, 1887–1894, `doi`. MR1230768

[130] R. Stanley, *A baker's dozen of conjectures concerning plane partitions*, Combinatoire énumérative (Montreal, Que., 1985/Quebec, Que., 1985), Lecture Notes in Math., vol. 1234, Springer, Berlin, 1986, pp. 285–293, `doi`. MR927770

[131] ———, *Symmetries of plane partitions*, J. Combin. Theory Ser. A **43** (1986), no. 1, 103–113, `doi`. MR859302

[132] ———, *Enumerative combinatorics. Volume 1*, second ed., Cambridge Studies in Advanced Mathematics, vol. 49, Cambridge University Press, Cambridge, 2012. MR2868112

[133] J. Stembridge, *Nonintersecting paths, Pfaffians, and plane partitions*, Adv. Math. **83** (1990), no. 1, 96–131, `doi`. MR1069389

[134] J. Striker, *The poset perspective on alternating sign matrices*, 21st International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2009), Discrete Math. Theor. Comput. Sci. Proc., AK, Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2009, pp. 813–824, `arXiv:0905.4495`. MR2721564

[135] ———, *A direct bijection between descending plane partitions with no special parts and permutation matrices*, Discrete Math. **311** (2011), no. 21, 2581–2585, `arXiv:1002.3391`, `doi`. MR2832158

[136] ———, *A unifying poset perspective on alternating sign matrices, plane partitions, Catalan objects, tournaments, and tableaux*, Adv. in Appl. Math. **46** (2011), no. 1-4, 583–609, `doi`. MR2794039

[137] Y. Stroganov, *The importance of being odd*, J. Phys. A **34** (2001), no. 13, L179–L185, `arXiv:cond-mat/0012035`, `doi`. MR1831119

[138] ———, *3-enumerated alternating sign matrices*, 2003, preprint, `arXiv:math-ph/0304004`.

[139] ———, *Izergin–Korepin determinant reloaded*, 2004, preprint, `arXiv:math-ph/0409072`.

[140] ———, *Izergin–Korepin determinant at a third root of unity*, Theoret. and Math. Phys. **146** (2006), no. 1, 53–62, `arXiv:math-ph/0204042`, `doi`. MR2243403

[141] ———, *$1/N$ phenomenon for some symmetry classes of the odd alternating sign matrices*, 2008, preprint, `arXiv:0807.2520`.

[142] T. Tokuyama, *A generating function of strict Gelfand patterns and some formulas on characters of general linear groups*, J. Math. Soc. Japan **40** (1988), no. 4, 671–685, `doi`. MR959093

[143] H. Turnbull, *The theory of determinants, matrices, and invariants*, 3rd ed, Dover Publications Inc., New York, 1960. MR0130257

[144] N. Williams, *An alternating sum of alternating sign matrices*, Rose Hulman Undergrad. Math. J. **9** (2008), no. 2, 6 pp. (electronic).

[145] D. Zeilberger, *Proof of the alternating sign matrix conjecture*, Electron. J. Combin. **3** (1996), no. 2, Research Paper 13, 84 pp. (electronic), The Foata Festschrift, `arXiv:math/9407211`. MR1392498

[146] ———, *Proof of the refined alternating sign matrix conjecture*, New York J. Math. **2** (1996), 59–68 (electronic), `arXiv:math/9606224`. MR1383799

[147] ———, *Dodgson's determinant-evaluation rule proved by two-timing men and women*, Electron. J. Combin. **4** (1997), no. 2, Research Paper 22, 2 pp. (electronic), The Wilf Festschrift (Philadelphia, PA, 1996), `arXiv:math/9808079`. MR1444169

[148] ———, *Dave Robbins' art of guessing*, Adv. in Appl. Math. **34** (2005), no. 4, 939–954, `doi`. MR2129005

[149] P. Zinn-Justin, *Six-vertex, loop and tiling models: integrability and combinatorics*, Lambert Academic Publishing, 2009, Habilitation thesis, `arXiv:0901.0665`.

[150] P. Zinn-Justin and P. Di Francesco, *Quantum Knizhnik-Zamolodchikov equation, totally symmetric self-complementary plane partitions, and alternating-sign matrices*, Theoret. and Math. Phys. **154** (2008), no. 3, 331–348, `arXiv:math-ph/0703015`, `doi`. MR2431554

R. E. BEHREND, SCHOOL OF MATHEMATICS, CARDIFF UNIVERSITY, CARDIFF, CF24 4AG, UK
E-mail address: `behrendr@cardiff.ac.uk`